

PYCRL Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21393c

Specification

PYCRL Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q53H96
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	28663

PYCRL Antibody (Center) - Additional Information

Gene ID 65263

Other Names

Pyrraline-5-carboxylate reductase 3, P5C reductase 3, P5CR 3, Pyrroline-5-carboxylate reductase-like protein, PYCRL

Target/Specificity

This PYCRL antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 139-171 amino acids from the Central region of human PYCRL.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PYCRL Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PYCRL Antibody (Center) - Protein Information

Name PYCR3 ([HGNC:25846](#))

Function Enzyme that catalyzes the last step in proline biosynthesis. Proline is synthesized from either glutamate or ornithine; both are converted to pyrroline-5-carboxylate (P5C), and then to proline via pyrroline-5-carboxylate reductases (PYCRs). PYCRL is exclusively linked to the

conversion of ornithine to proline.

Cellular Location


Cytoplasm.

PYCRL Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PYCRL Antibody (Center) - Images



Anti-PYCRL Antibody (Center) at 1:2000 dilution + KG-1 whole cell lysates. Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 29 kDa. Blocking/Dilution buffer: 5% NFD/MTBST.

PYCRL Antibody (Center) - References

- Ota T., et al. *Nat. Genet.* 36:40-45(2004).
Nusbaum C., et al. *Nature* 439:331-335(2006).
Suzuki Y., et al. Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Bechtel S., et al. *BMC Genomics* 8:399-399(2007).
Burkard T.R., et al. *BMC Syst. Biol.* 5:17-17(2011).