

**DBH Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21423c**

**Specification**

---

**DBH Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P09172</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	69065

**DBH Antibody (Center) - Additional Information**

**Gene ID** 1621

**Other Names**

Dopamine beta-hydroxylase, Dopamine beta-monoxygenase, Soluble dopamine beta-hydroxylase, DBH

**Target/Specificity**

This DBH antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 270-303 amino acids from the Central region of human DBH.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

DBH Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**DBH Antibody (Center) - Protein Information**

**Name** DBH

**Function** Catalyzes the hydroxylation of dopamine to noradrenaline (also known as norepinephrine), and is thus vital for regulation of these neurotransmitters.

### Cellular Location


[Soluble dopamine beta-hydroxylase]: Cytoplasmic vesicle, secretory vesicle lumen Cytoplasmic vesicle, secretory vesicle, chromaffin granule lumen. Secreted

### DBH Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### DBH Antibody (Center) - Images



All lanes : Anti-DBH Antibody (Center) at 1:2000 dilution Lane 1: SH-SY5Y whole cell lysates Lane 2: human plasma lysates Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 69 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### DBH Antibody (Center) - Background

Conversion of dopamine to noradrenaline.

### DBH Antibody (Center) - References

- Humphray S.J., et al. Nature 429:369-374(2004).  
Lamouroux A., et al. EMBO J. 6:3931-3937(1987).  
Kobayashi K., et al. Nucleic Acids Res. 17:1089-1102(1989).  
Li B., et al. Biochem. J. 313:57-64(1996).  
Liu T., et al. J. Proteome Res. 4:2070-2080(2005).