

OXTR Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21446b

Specification

OXTR Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P30559
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	42772

OXTR Antibody (C-term) - Additional Information

Gene ID 5021

Other Names

Oxytocin receptor, OT-R, OXTR

Target/Specificity

This OXTR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 348-380 amino acids from the C-terminal region of human OXTR.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OXTR Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OXTR Antibody (C-term) - Protein Information

Name OXTR

Function Receptor for oxytocin. The activity of this receptor is mediated by G proteins which activate a phosphatidylinositol-calcium second messenger system.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OXTR Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OXTR Antibody (C-term) - Images



Anti-OXTR Antibody (C-term) at 1:1000 dilution + Jurkat whole cell lysates. Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 43 kDa. Blocking/Dilution buffer: 5% NFD/MBST.

OXTR Antibody (C-term) - Background

Receptor for oxytocin. The activity of this receptor is mediated by G proteins which activate a phosphatidylinositol- calcium second messenger system.

OXTR Antibody (C-term) - References

- Kimura T., et al. Nature 356:526-529(1992).
Kimura T., et al. Nature 357:176-176(1992).
Kopatz S.A., et al. Submitted (SEP-2003) to the EMBL/GenBank/DDBJ databases.
Inoue T., et al. J. Biol. Chem. 269:32451-32456(1994).