

AKR7L Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21497a

Specification

AKR7L Antibody (N-Term) - Product Information

Application	IF, WB,E
Primary Accession	Q8NHP1
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	36970
Antigen Region	32-64

AKR7L Antibody (N-Term) - Additional Information

Gene ID 246181

Other Names

Aflatoxin B1 aldehyde reductase member 4, 1---, AFB1 aldehyde reductase 3, AFB1-AR 3, Aldoketoreductase 7-like, AKR7L, AFAR3, AKR7A4

Target/Specificity

This AKR7L antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 32-64 amino acids from human AKR7L.

Dilution

IF~~1:25

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AKR7L Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

AKR7L Antibody (N-Term) - Protein Information

Name AKR7L

Synonyms AFAR3 {ECO:0000303|PubMed:12879023}, AKR

Function Can reduce the dialdehyde protein-binding form of aflatoxin B1 (AFB1) to the non-binding AFB1 dialcohol. May be involved in protection of liver against the toxic and carcinogenic effects of AFB1, a potent hepatocarcinogen (By similarity).

Tissue Location


Mainly expressed in uterus.

AKR7L Antibody (N-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AKR7L Antibody (N-Term) - Images



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HepG2 (human liver hepatocellular carcinoma cell line) cells labeling AKR7L with AP21497a at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (NK179883) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on HepG2 cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red).The nuclear counter stain is DAPI (blue).



Anti-AKR7L Antibody (N-Term) at 1:2000 dilution + HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

AKR7L Antibody (N-Term) - Background

Can reduce the dialdehyde protein-binding form of aflatoxin B1 (AFB1) to the non-binding AFB1 dialcohol. May be involved in protection of liver against the toxic and carcinogenic effects of AFB1, a potent hepatocarcinogen (By similarity).

AKR7L Antibody (N-Term) - References

Gregory S.G., et al. Nature 441:315-321(2006).
Prabl C., et al. Oncogene 22:4765-4773(2003).