

**SULT1A1 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21514c**

**Specification**

---

**SULT1A1 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P50225</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG

**SULT1A1 Antibody (Center) - Additional Information**

**Gene ID** 6817

**Other Names**

Sulfotransferase 1A1, ST1A1, Aryl sulfotransferase 1, HAST1/HAST2, Phenol sulfotransferase 1, Phenol-sulfating phenol sulfotransferase 1, P-PST 1, ST1A3, Thermostable phenol sulfotransferase, Ts-PST, SULT1A1, STP, STP1

**Target/Specificity**

This SULT1A1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 90-123 amino acids of human SULT1A1.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

SULT1A1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**SULT1A1 Antibody (Center) - Protein Information**

**Name** SULT1A1

**Synonyms** STP, STP1

**Function** Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to

catalyze the sulfate conjugation of a wide variety of acceptor molecules bearing a hydroxyl or an amine group. Sulfonation increases the water solubility of most compounds, and therefore their renal excretion, but it can also result in bioactivation to form active metabolites. Displays broad substrate specificity for small phenolic compounds. Plays an important role in the sulfonation of endogenous molecules such as steroid hormones and 3,3'-diiodothyronin (PubMed:[10199779](#), PubMed:[12471039](#), PubMed:[16221673](#), PubMed:[21723874](#), PubMed:[22069470](#), PubMed:[7834621](#)). Mediates the sulfate conjugation of a variety of xenobiotics, including the drugs acetaminophen and minoxidil (By similarity). Mediates also the metabolic activation of carcinogenic N-hydroxyarylamines leading to highly reactive intermediates capable of forming DNA adducts, potentially resulting in mutagenesis (PubMed:[7834621](#)). May play a role in gut microbiota-host metabolic interaction. O-sulfonates 4-ethylphenol (4-EP), a dietary tyrosine-derived metabolite produced by gut bacteria. The product 4-EPS crosses the blood-brain barrier and may negatively regulate oligodendrocyte maturation and myelination, affecting the functional connectivity of different brain regions associated with the limbic system.

#### Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P17988}.

#### Tissue Location


Liver, lung, adrenal, brain, platelets and skin.

### SULT1A1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### SULT1A1 Antibody (Center) - Images



Anti-SULT1A1 Antibody (Center) at 1:2000 dilution + human lung lysates Lysates/proteins at 20

µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution  
Predicted band size : 34 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

### **SULT1A1 Antibody (Center) - Background**

Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the sulfate conjugation of catecholamines, phenolic drugs and neurotransmitters. Has also estrogen sulfotransferase activity. responsible for the sulfonation and activation of minoxidil. Is Mediates the metabolic activation of carcinogenic N- hydroxyarylamines to DNA binding products and could so participate as modulating factor of cancer risk.

### **SULT1A1 Antibody (Center) - References**

- Zhu X.,et al.Biochem. Biophys. Res. Commun. 195:120-127(1993).  
Zhu X.,et al.Biochem. Biophys. Res. Commun. 192:671-676(1993).  
Wilborn T.W.,et al.Mol. Pharmacol. 43:70-77(1993).  
Yamazoe Y.,et al.Chem. Biol. Interact. 92:107-117(1994).  
Hwang S.-R.,et al.Biochem. Biophys. Res. Commun. 207:701-707(1995).