

HNRNPUL2 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21566c

Specification

HNRNPUL2 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q1KMD3
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	85105

HNRNPUL2 Antibody (Center) - Additional Information

Gene ID 221092

Other Names

Heterogeneous nuclear ribonucleoprotein U-like protein 2, Scaffold-attachment factor A2, SAF-A2, HNRNPUL2, HNRPUL2

Target/Specificity

This HNRNPUL2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 337-350 amino acids from the Central region of human HNRNPUL2.

Dilution

WB~~1:2000

IHC-P~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HNRNPUL2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

HNRNPUL2 Antibody (Center) - Protein Information

Name HNRNPUL2

Synonyms HNRPUL2

Cellular Location


Nucleus {ECO:0000250|UniProtKB:Q00PI9}.

HNRNPUL2 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


HNRNPUL2 Antibody (Center) - Images



Anti-HNRNPUL2 Antibody (Center) at 1:8000 dilution + HeLa whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 85 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



All lanes : Anti-HNRNPUL2 Antibody (Center) at 1:2000 dilution Lane 1: MCF-7 whole cell lysates Lane 2: Jurkat whole cell lysates Lane 3: 293T/17 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 85 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP21566c staining HNRNPUL2 in human stomach tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

HNRNPUL2 Antibody (Center) - References

- Herrmann F., et al. Submitted (MAR-2006) to the EMBL/GenBank/DDBJ databases.
- Bechtel S., et al. BMC Genomics 8:399-399(2007).
- Bienvenut W.V., et al. Submitted (DEC-2008) to UniProtKB.
- Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).