

**PSMB3 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21598c**

## Specification

---

### PSMB3 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	<a href="#">P49720</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	22949

### PSMB3 Antibody (Center) - Additional Information

**Gene ID** 5691

#### Other Names

Proteasome subunit beta type-3, Proteasome chain 13, Proteasome component C10-II, Proteasome theta chain, PSMB3

#### Target/Specificity

This PSMB3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 150-185 amino acids from the Central region of human PSMB3.

#### Dilution

WB~~1:2000

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

PSMB3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### PSMB3 Antibody (Center) - Protein Information

**Name** PSMB3

**Function** Non-catalytic component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S

regulatory particles, forms the 26S proteasome and thus participates in the ATP-dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex).

#### Cellular Location


Cytoplasm. Nucleus. Note=Translocated from the cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import receptor IPO9

#### PSMB3 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### PSMB3 Antibody (Center) - Images



All lanes : Anti-PSMB3 Antibody (Center) at 1:2000 dilution Lane 1: K562 whole cell lysates Lane 2: HeLa whole cell lysates Lane 3: Jurkat whole cell lysates Lane 4: 293T/17 whole cell lysates Lane 5: NIH/3T3 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 23 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

#### PSMB3 Antibody (Center) - Background

The proteasome is a multicatalytic proteinase complex which is characterized by its ability to

cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity.

### **PSMB3 Antibody (Center) - References**

- Nothwang H.G.,et al.Biochim. Biophys. Acta 1219:361-368(1994).  
Bienvenut W.V.,et al.Submitted (JUL-2008) to UniProtKB.  
Lubec G.,et al.Submitted (MAR-2007) to UniProtKB.  
Rasmussen H.H.,et al.Electrophoresis 13:960-969(1992).  
Kristensen P.,et al.Biochem. Biophys. Res. Commun. 205:1785-1789(1994).