

RAE1 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21654a**Specification**

RAE1 Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	P78406
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	40968

RAE1 Antibody (N-Term) - Additional Information**Gene ID** 8480**Other Names**

mRNA export factor, Rae1 protein homolog, mRNA-associated protein mrnp 41, RAE1, MRNP41

Target/Specificity

This RAE1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 19-42 amino acids from human RAE1.

Dilution

WB~~1:2000-1:4000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAE1 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

RAE1 Antibody (N-Term) - Protein Information**Name** RAE1**Synonyms** MRNP41**Function** Acts as a mRNA export factor involved in nucleocytoplasmic transport (PubMed:[20498086](#), PubMed:[33849972](#)). Plays a role in mitotic bipolar spindle formation

(PubMed:[17172455](#)). May function in attaching cytoplasmic mRNPs to the cytoskeleton both directly or indirectly (PubMed:[17172455](#)).

Cellular Location


Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, spindle pole. Nucleus envelope. Note=Recruited from interphase nuclei to spindle MTs during mitosis.

RAE1 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAE1 Antibody (N-Term) - Images



All lanes : Anti-RAE1 Antibody (N-Term) at 1:2000-1:4000 dilution Lane 1: HeLa whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: MCF-7 whole cell lysate Lane 5: SH-SY5Y whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 41 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

RAE1 Antibody (N-Term) - Background

Binds mRNA. May function in nucleocytoplasmic transport and in directly or indirectly attaching cytoplasmic mRNPs to the cytoskeleton.

RAE1 Antibody (N-Term) - References

Bharathi A.,et al.Gene 198:251-258(1997).
Kraemer D.M.,et al.Proc. Natl. Acad. Sci. U.S.A. 94:9119-9124(1997).

Lu L.,et al.Submitted (JUL-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Deloukas P.,et al.Nature 414:865-871(2001).