

Mouse Ptk7 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21668b

Specification

Mouse Ptk7 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q8BKG3
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	117532

Mouse Ptk7 Antibody (C-term) - Additional Information

Gene ID 71461

Other Names

Inactive tyrosine-protein kinase 7, Protein chuzhoi, Protein-tyrosine kinase 7, Pseudo tyrosine kinase receptor 7, Tyrosine-protein kinase-like 7, Ptk7

Target/Specificity

This Mouse Ptk7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 702-737 amino acids from the C-terminal region of Mouse Ptk7.

Dilution

WB~~1:1000-1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Ptk7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Ptk7 Antibody (C-term) - Protein Information

Name Ptk7

Function Inactive tyrosine kinase involved in Wnt signaling pathway. Component of both the non-canonical (also known as the Wnt/planar cell polarity signaling) and the canonical Wnt signaling pathway. Functions in cell adhesion, cell migration, cell polarity, proliferation, actin

cytoskeleton reorganization and apoptosis. Has a role in embryogenesis, epithelial tissue organization and angiogenesis.

Cellular Location

Membrane; Single-pass type I membrane protein. Cell junction. Note=Colocalizes with MMP14 at cell junctions. Also localizes at the leading edge of migrating cells

Tissue Location


Expressed at high levels in lung and un-pregnant uterus among adult tissues, and in the tail, limbs, somites, gut and craniofacial regions among embryonic tissues

Mouse Ptk7 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Mouse Ptk7 Antibody (C-term) - Images



Anti-Ptk7 Antibody (C-term) at 1:1000 dilution + C2C12 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 118 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-Ptk7 Antibody (C-term) at 1:2000 dilution + NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 118 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-Ptk7 Antibody (C-term) at 1:1000-1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: mouse lung lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 118 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Mouse Ptk7 Antibody (C-term) - Background

Inactive tyrosine kinase involved in Wnt signaling pathway. Component of both the non-canonical (also known as the Wnt/planar cell polarity signaling) and the canonical Wnt signaling pathway. Functions in cell adhesion, cell migration, cell polarity, proliferation, actin cytoskeleton reorganization and apoptosis. Has a role in embryogenesis, epithelial tissue organization and angiogenesis.

Mouse Ptk7 Antibody (C-term) - References

- Jung J.-W., et al. Gene 328:75-84(2004).
- Carninci P., et al. Science 309:1559-1563(2005).
- Daigo Y., et al. Submitted (FEB-2001) to the EMBL/GenBank/DDBJ databases.

Lu X., et al. Nature 430:93-98(2004).

Gundry R.L., et al. Mol. Cell. Proteomics 8:2555-2569(2009).