

CYP27B1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21674b

Specification

CYP27B1 Antibody (C-term) - Product Information

| | |
|-------------------|------------------------|
| Application | WB,E |
| Primary Accession | O15528 |
| Reactivity | Mouse |
| Host | Rabbit |
| Clonality | polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 56504 |

CYP27B1 Antibody (C-term) - Additional Information

Gene ID 1594

Other Names

25-hydroxyvitamin D-1 alpha hydroxylase, mitochondrial, 25-OHD-1 alpha-hydroxylase, 25-hydroxyvitamin D(3) 1-alpha-hydroxylase, VD3 1A hydroxylase, Calcdiol 1-monooxygenase, Cytochrome P450 subfamily XXVIIIB polypeptide 1, Cytochrome P450C1 alpha, Cytochrome P450VD1-alpha, Cytochrome p450 27B1, CYP27B1, CYP1ALPHA, CYP27B

Target/Specificity

This CYP27B1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 400-429 amino acids from the C-terminal region of human CYP27B1.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CYP27B1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CYP27B1 Antibody (C-term) - Protein Information

Name CYP27B1

Synonyms CYP1ALPHA, CYP27B

Function A cytochrome P450 monooxygenase involved in vitamin D metabolism and in calcium and phosphorus homeostasis. Catalyzes the rate-limiting step in the activation of vitamin D in the kidney, namely the hydroxylation of 25-hydroxyvitamin D3/calcidiol at the C1 α - position to form the hormonally active form of vitamin D3, 1 α ,25- dihydroxyvitamin D3/calcitriol that acts via the vitamin D receptor (VDR) (PubMed:[10518789](#), PubMed:[10566658](#), PubMed:[12050193](#), PubMed:[22862690](#), PubMed:[9486994](#)). Has 1 α -hydroxylase activity on vitamin D intermediates of the CYP24A1-mediated inactivation pathway (PubMed:[10518789](#), PubMed:[22862690](#)). Converts 24R,25-dihydroxyvitamin D3/secalciferol to 1- α ,24,25-trihydroxyvitamin D3, an active ligand of VDR. Also active on 25-hydroxyvitamin D2 (PubMed:[10518789](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via FDXR/adrenodoxin reductase and FDX1/adrenodoxin (PubMed:[22862690](#)).

Cellular Location

Mitochondrion membrane.

Tissue Location


Kidney.

CYP27B1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CYP27B1 Antibody (C-term) - Images



Anti-CYP27B1 Antibody (C-term) at 1:2000 dilution + mouse kidney lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 56 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

CYP27B1 Antibody (C-term) - Background

Catalyzes the conversion of 25-hydroxyvitamin D3 (25(OH)D) to 1-alpha,25-dihydroxyvitamin D3 (1,25(OH)2D) plays an important role in normal bone growth, calcium metabolism, and tissue differentiation.

CYP27B1 Antibody (C-term) - References

Fu G.K.,et al.DNA Cell Biol. 16:1499-1507(1997).
Monkawa T.,et al.Biochem. Biophys. Res. Commun. 239:527-533(1997).
Fu G.K.,et al.Mol. Endocrinol. 11:1961-1970(1997).
Huang D.C.,et al.Mol. Cancer Res. 1:56-67(2002).
Huang D.C.,et al.Submitted (MAR-2000) to the EMBL/GenBank/DDBJ databases.