

**COX6C Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21744b**

**Specification**

---

**COX6C Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P09669</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	8781

**COX6C Antibody (C-term) - Additional Information**

**Gene ID** 1345

**Other Names**

Cytochrome c oxidase subunit 6C, Cytochrome c oxidase polypeptide VIc, COX6C

**Target/Specificity**

This COX6C antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 41-75 amino acids from the C-terminal region of human COX6C.

**Dilution**

WB~~1:8000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

COX6C Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**COX6C Antibody (C-term) - Protein Information**

**Name** COX6C

**Function** Component of the cytochrome c oxidase, the last enzyme in the mitochondrial electron transport chain which drives oxidative phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII), ubiquinol- cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex

IV, CIV), that cooperate to transfer electrons derived from NADH and succinate to molecular oxygen, creating an electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase. Cytochrome c oxidase is the component of the respiratory chain that catalyzes the reduction of oxygen to water. Electrons originating from reduced cytochrome c in the intermembrane space (IMS) are transferred via the dinuclear copper A center (CU(A)) of subunit 2 and heme A of subunit 1 to the active site in subunit 1, a binuclear center (BNC) formed by heme A3 and copper B (CU(B)). The BNC reduces molecular oxygen to 2 water molecules using 4 electrons from cytochrome c in the IMS and 4 protons from the mitochondrial matrix.

#### Cellular Location


Mitochondrion inner membrane; Single-pass membrane protein

#### COX6C Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### COX6C Antibody (C-term) - Images



All lanes : Anti-COX6C Antibody (C-term) at 1:8000 dilution Lane 1: human heart lysate Lane 2: A-673 whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: U-2OS whole cell lysate Lane 5: mouse heart lysate Lane 6: rat kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 9 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

#### COX6C Antibody (C-term) - Background

This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal

oxidase in mitochondrial electron transport.

#### **COX6C Antibody (C-term) - References**

- Otsuka M.,et al.Nucleic Acids Res. 16:10916-10916(1988).  
Ohta S.,et al.Submitted (FEB-1996) to the EMBL/GenBank/DDBJ databases.  
Hofmann S.,et al.Cytogenet. Cell Genet. 83:226-227(1998).  
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).