

**SEMA7A Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21852b**

**Specification**

---

**SEMA7A Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O75326</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	74824

**SEMA7A Antibody (C-term) - Additional Information**

**Gene ID** 8482

**Other Names**

Semaphorin-7A, CDw108, JMH blood group antigen, John-Milton-Hargen human blood group Ag, Semaphorin-K1, Sema K1, Semaphorin-L, Sema L, CD108, SEMA7A, CD108, SEMAL

**Target/Specificity**

This SEMA7A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 521-555 amino acids from the C-terminal region of human SEMA7A.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

SEMA7A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**SEMA7A Antibody (C-term) - Protein Information**

**Name** SEMA7A

**Synonyms** CD108, SEMAL

**Function** Plays an important role in integrin-mediated signaling and functions both in regulating

cell migration and immune responses. Promotes formation of focal adhesion complexes, activation of the protein kinase PTK2/FAK1 and subsequent phosphorylation of MAPK1 and MAPK3. Promotes production of pro-inflammatory cytokines by monocytes and macrophages. Plays an important role in modulating inflammation and T-cell-mediated immune responses. Promotes axon growth in the embryonic olfactory bulb. Promotes attachment, spreading and dendrite outgrowth in melanocytes.

#### Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor; Extracellular side. Note=Detected in a punctate pattern on the cell membrane of basal and supra-basal skin keratinocytes

#### Tissue Location


Detected in skin keratinocytes and on endothelial cells from skin blood vessels (at protein level). Expressed in fibroblasts, keratinocytes, melanocytes, placenta, testis, ovary, spleen, brain, spinal chord, lung, heart, adrenal gland, lymph nodes, thymus, intestine and kidney.

### SEMA7A Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### SEMA7A Antibody (C-term) - Images



All lanes : Anti-SEMA7A Antibody (C-term) at 1:2000 dilution Lane 1: mouse brain lysate Lane 2: A2058 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 75 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

### SEMA7A Antibody (C-term) - Background

Plays an important role in integrin-mediated signaling and functions both in regulating cell migration and immune responses. Promotes formation of focal adhesion complexes, activation of the protein kinase PTK2/FAK1 and subsequent phosphorylation of MAPK1 and MAPK3. Promotes production of proinflammatory cytokines by monocytes and macrophages. Plays an important role in modulating inflammation and T-cell-mediated immune responses. Promotes axon growth in the embryonic olfactory bulb. Promotes attachment, spreading and dendrite outgrowth in melanocytes.

#### **SEMA7A Antibody (C-term) - References**

Lange C., et al. Genomics 51:340-350(1998).  
Yamada A., et al. J. Immunol. 162:4094-4100(1999).  
Xu X., et al. J. Biol. Chem. 273:22428-22434(1998).  
Seltsam A., et al. Transfusion 47:133-146(2007).  
Ota T., et al. Nat. Genet. 36:40-45(2004).