

SEMA7A Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21852b

Specification

SEMA7A Antibody (C-term) - Product Information

| | |
|-------------------|------------------------|
| Application | WB,E |
| Primary Accession | 075326 |
| Reactivity | Human, Mouse |
| Host | Rabbit |
| Clonality | polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 74824 |

SEMA7A Antibody (C-term) - Additional Information

Gene ID 8482

Other Names

Semaphorin-7A, CDw108, JMH blood group antigen, John-Milton-Hargen human blood group Ag, Semaphorin-K1, Sema K1, Semaphorin-L, Sema L, CD108, SEMA7A, CD108, SEMAL

Target/Specificity

This SEMA7A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 521-555 amino acids from the C-terminal region of human SEMA7A.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SEMA7A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SEMA7A Antibody (C-term) - Protein Information

Name SEMA7A

Synonyms CD108, SEMAL

Function Plays an important role in integrin-mediated signaling and functions both in regulating

cell migration and immune responses. Promotes formation of focal adhesion complexes, activation of the protein kinase PTK2/FAK1 and subsequent phosphorylation of MAPK1 and MAPK3. Promotes production of pro-inflammatory cytokines by monocytes and macrophages. Plays an important role in modulating inflammation and T-cell-mediated immune responses. Promotes axon growth in the embryonic olfactory bulb. Promotes attachment, spreading and dendrite outgrowth in melanocytes.

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor; Extracellular side. Note=Detected in a punctate pattern on the cell membrane of basal and supra-basal skin keratinocytes

Tissue Location


Detected in skin keratinocytes and on endothelial cells from skin blood vessels (at protein level). Expressed in fibroblasts, keratinocytes, melanocytes, placenta, testis, ovary, spleen, brain, spinal chord, lung, heart, adrenal gland, lymph nodes, thymus, intestine and kidney.

SEMA7A Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SEMA7A Antibody (C-term) - Images



All lanes : Anti-SEMA7A Antibody (C-term) at 1:2000 dilution Lane 1: mouse brain lysate Lane 2: A2058 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 75 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

SEMA7A Antibody (C-term) - Background

Plays an important role in integrin-mediated signaling and functions both in regulating cell migration and immune responses. Promotes formation of focal adhesion complexes, activation of the protein kinase PTK2/FAK1 and subsequent phosphorylation of MAPK1 and MAPK3. Promotes production of proinflammatory cytokines by monocytes and macrophages. Plays an important role in modulating inflammation and T-cell-mediated immune responses. Promotes axon growth in the embryonic olfactory bulb. Promotes attachment, spreading and dendrite outgrowth in melanocytes.

SEMA7A Antibody (C-term) - References

Lange C., et al. Genomics 51:340-350(1998).
Yamada A., et al. J. Immunol. 162:4094-4100(1999).
Xu X., et al. J. Biol. Chem. 273:22428-22434(1998).
Seltman A., et al. Transfusion 47:133-146(2007).
Ota T., et al. Nat. Genet. 36:40-45(2004).