

MYBL2 Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21853b

Specification

MYBL2 Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	P10244
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	78764

MYBL2 Antibody (C-Term) - Additional Information

Gene ID 4605

Other Names

Myb-related protein B, B-Myb, Myb-like protein 2, MYBL2, BMYB

Target/Specificity

This MYBL2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 540-571 amino acids from human MYBL2.

Dilution

WB~~1:8000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MYBL2 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

MYBL2 Antibody (C-Term) - Protein Information

Name MYBL2

Synonyms BMYB

Function Transcription factor involved in the regulation of cell survival, proliferation, and differentiation. Transactivates the expression of the CLU gene.

Cellular Location


Nucleus.

MYBL2 Antibody (C-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MYBL2 Antibody (C-Term) - Images



All lanes : Anti-MYBL2 Antibody (C-Term) at 1:2000 dilution Lane 1: 293T/17 whole cell lysate Lane 2: K562 whole cell lysate Lane 3: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 79 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



All lanes : Anti-MYBL2 Antibody (C-Term) at 1:8000 dilution Lane 1: CEM whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: mouse testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 79 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

MYBL2 Antibody (C-Term) - Background

Transcription factor involved in the regulation of cell survival, proliferation, and differentiation. Transactivates the expression of the CLU gene.

MYBL2 Antibody (C-Term) - References

Nomura N.,et al.Nucleic Acids Res. 16:11075-11089(1988).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Totoki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Deloukas P.,et al.Nature 414:865-871(2001).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.