

PHC3 Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21862b

Specification

PHC3 Antibody (C-Term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q8NDX5
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	106162

PHC3 Antibody (C-Term) - Additional Information

Gene ID 80012

Other Names

Polyhomeotic-like protein 3, Early development regulatory protein 3, Homolog of polyhomeotic 3, hPH3, PHC3, EDR3, PH3

Target/Specificity

This PHC3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 819-850 amino acids from human PHC3.

Dilution

WB~~1:2000

IHC-P~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PHC3 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

PHC3 Antibody (C-Term) - Protein Information

Name PHC3

Synonyms EDR3, PH3

Function Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility.

Cellular Location


Nucleus

PHC3 Antibody (C-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PHC3 Antibody (C-Term) - Images



All lanes : Anti-PHC3 Antibody (C-Term) at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: SH-SY5Y whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: U-2OS whole cell lysate Lane 5: A549 whole cell lysate Lane 6: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 106 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



AP21862b staining PHC3 in human kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

PHC3 Antibody (C-Term) - Background

Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility.

PHC3 Antibody (C-Term) - References

Tonkin E., et al. Hum. Genet. 111:435-442(2002).
Levine S.S., et al. Mol. Cell. Biol. 22:6070-6078(2002).
Hansen M.F., et al. Submitted (MAY-2001) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).
Bechtel S., et al. BMC Genomics 8:399-399(2007).