

ST3GAL4 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22025a

Specification

ST3GAL4 Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	O11206
Other Accession	P61130
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	38045

ST3GAL4 Antibody (N-Term) - Additional Information

Gene ID 6484

Other Names

CMP-N-acetylneuraminase-beta-galactosamide-alpha-2, 3-sialyltransferase 4, Alpha 2, 3-ST 4, Beta-galactoside alpha-2, 3-sialyltransferase 4, 2.4.99.-, Alpha 2, 3-sialyltransferase IV, Gal-NAc6S, Gal-beta-1, 4-GalNAc-alpha-2, 3-sialyltransferase, SAT-3, ST-4, ST3Gal IV, ST3GalIV, ST3GalA.2, STZ, Sialyltransferase 4C, SIAT4-C, ST3GAL4, CGS23, NANTA3, SIAT4C, STZ

Target/Specificity

This ST3GAL4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 26-57 amino acids from human ST3GAL4.

Dilution

WB~~1:1000-1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ST3GAL4 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

ST3GAL4 Antibody (N-Term) - Protein Information

Name ST3GAL4

Synonyms CGS23, NANTA3, SIAT4C, STZ

Function A beta-galactoside alpha2-3 sialyltransferase involved in terminal sialylation of glycoproteins and glycolipids (PubMed:[8288606](#), PubMed:[8611500](#)). Catalyzes the transfer of sialic acid (N-acetyl- neuraminic acid; Neu5Ac) from the nucleotide sugar donor CMP-Neu5Ac onto acceptor Galbeta-(1->3)-GalNAc- and Galbeta-(1->4)-GlcNAc- terminated glycoconjugates through an alpha2-3 linkage (PubMed:[8288606](#), PubMed:[8611500](#)). Plays a major role in hemostasis. Responsible for sialylation of plasma VWF/von Willebrand factor, preventing its recognition by asialoglycoprotein receptors (ASGPR) and subsequent clearance. Regulates ASGPR-mediated clearance of platelets (By similarity). Participates in the biosynthesis of the sialyl Lewis X epitopes, both on O- and N-glycans, which are recognized by SELE/E- selectin, SELP/P-selectin and SELL/L-selectin. Essential for selectin- mediated rolling and adhesion of leukocytes during extravasation (PubMed:[25498912](#)). Contributes to adhesion and transendothelial migration of neutrophils likely through terminal sialylation of CXCR2 (By similarity). In glycosphingolipid biosynthesis, sialylates GM1 and GA1 gangliosides to form GD1a and GM1b, respectively (PubMed:[8288606](#)). Metabolizes brain c-series ganglioside GT1c forming GQ1c (By similarity). Synthesizes ganglioside LM1 (IV3Neu5Ac-nLc4Cer), a major structural component of peripheral nerve myelin (PubMed:[8611500](#)).

Cellular Location

Golgi apparatus, Golgi stack membrane; Single- pass type II membrane protein. Secreted. Note=Membrane-bound form in trans cisternae of Golgi. Secreted into the body fluid

Tissue Location

Highly expressed in adult placenta, heart and kidney.


ST3GAL4 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ST3GAL4 Antibody (N-Term) - Images





All lanes : Anti-ST3GAL4 Antibody (N-Term) at 1:1000-1:2000 dilution Lane 1: PC-3 whole cell lysate Lane 2: SK-OV-3 whole cell lysate Lane 3: HeLa whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

ST3GAL4 Antibody (N-Term) - Background

It may catalyze the formation of the NeuAc-alpha-2,3- Gal-beta-1,3-GalNAc- or NeuAc-alpha-2,3-Gal-beta-1,3-GlcNAc- sequences found in terminal carbohydrate groups of glycoproteins and glycolipids. It may be involved in the biosynthesis of the sialyl Lewis X determinant. Also acts on the corresponding 1,3- galactosyl derivative.

ST3GAL4 Antibody (N-Term) - References

Kitagawa H.,et al.J. Biol. Chem. 271:931-938(1996).
Kitagawa H.,et al.J. Biol. Chem. 269:1394-1401(1994).
Sasaki K.,et al.J. Biol. Chem. 268:22782-22787(1993).
Grahm A.,et al.Glycoconj. J. 18:759-767(2001).
Ota T.,et al.Nat. Genet. 36:40-45(2004).