

**NT5C Antibody (N-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP22072a****Specification**

---

**NT5C Antibody (N-Term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q8TCD5</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	23383

**NT5C Antibody (N-Term) - Additional Information****Gene ID** 30833**Other Names**

5'(3')-deoxyribonucleotidase, cytosolic type, 3.1.3.-, Cytosolic 5', 3'-pyrimidine nucleotidase, Deoxy-5'-nucleotidase 1, dNT-1, NT5C, DNT1, UMPH2

**Target/Specificity**

This NT5C antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 27-60 amino acids from human NT5C.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NT5C Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**NT5C Antibody (N-Term) - Protein Information****Name** NT5C**Synonyms** DNT1, UMPH2**Function** Dephosphorylates the 5' and 2'(3')-phosphates of deoxyribonucleotides, with a

preference for dUMP and dTMP, intermediate activity towards dGMP, and low activity towards dCMP and dAMP.

#### Cellular Location

Cytoplasm.

#### Tissue Location


Detected in skeletal muscle, heart and pancreas.

### NT5C Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### NT5C Antibody (N-Term) - Images



All lanes : Anti-NT5C Antibody (N-Term) at 1:2000 dilution Lane 1: Raji whole cell lysate Lane 2: U-251 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 23 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### NT5C Antibody (N-Term) - Background

Dephosphorylates the 5' and 2'(3')-phosphates of deoxyribonucleotides, with a preference for dUMP and dTMP, intermediate activity towards dGMP, and low activity towards dCMP and dAMP.

### NT5C Antibody (N-Term) - References

Rampazzo C., et al. J. Biol. Chem. 275:5409-5415(2000).

Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Zody M.C.,et al.Nature 440:1045-1049(2006).  
Rampazzo C.,et al.Gene 294:109-117(2002).  
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).