

Vinculin Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22112a

Specification

Vinculin Antibody - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q64727
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	116717

Vinculin Antibody - Additional Information

Gene ID 22330

Other Names

Vinculin, Metavinculin, Vcl

Target/Specificity

This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 634-668 amino acids from mouse.

Dilution

WB~~1:2000

IHC-P~~1:25

FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Vinculin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Vinculin Antibody - Protein Information

Name Vcl

Function Actin filament (F-actin)-binding protein involved in cell- matrix adhesion and cell-cell adhesion. Regulates cell-surface E- cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play important roles in cell morphology and locomotion (By

similarity).

Cellular Location


Cell membrane {ECO:0000250|UniProtKB:P12003}; Peripheral membrane protein {ECO:0000250|UniProtKB:P12003}; Cytoplasmic side {ECO:0000250|UniProtKB:P12003}. Cell junction, adherens junction {ECO:0000250|UniProtKB:P12003}. Cell junction, focal adhesion {ECO:0000250|UniProtKB:P12003}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P85972}. Cell membrane, sarcolemma; Peripheral membrane protein; Cytoplasmic side. Cell projection, podosome. Note=Recruitment to cell-cell junctions occurs in a myosin II-dependent manner. Interaction with CTNNB1 is necessary for its localization to the cell-cell junctions {ECO:0000250|UniProtKB:P12003}

Vinculin Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Vinculin Antibody - Images




All lanes : Anti-Vinculin at 1:2000 dilution Lane 1: human skeletal muscle lysate Lane 2: A431 whole cell lysate Lane 3: HeLa whole cell lysate Lane 4: mouse kidney lysate Lane 5: PC-12 whole cell lysate Lane 6: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 117 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP22112a staining Vinculin in mouse skeletal muscle tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing NIH/3T3 cells stained with AP22112a (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22112a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.



Overlay histogram showing C2C12 cells stained with AP22112a (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22112a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed (OH191631) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

Vinculin Antibody - Background

Actin filament (F-actin)-binding protein involved in cell-matrix adhesion and cell-cell adhesion. Regulates cell-surface E-cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play important roles in cell morphology and locomotion (By similarity).

Vinculin Antibody - References

Coll J.-L., et al. Proc. Natl. Acad. Sci. U.S.A. 92:9161-9165 (1995).
 Alatorsev V.E., et al. FEBS Lett. 413:197-201 (1997).
 Carninci P., et al. Science 309:1559-1563 (2005).
 Lubec G., et al. Submitted (JAN-2009) to UniProtKB.
 Mandai K., et al. J. Cell Biol. 144:1001-1017 (1999).