

**SLC2A3 Antibody (C-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP22174b**

**Specification**

---

**SLC2A3 Antibody (C-Term) - Product Information**

Application	WB, FC,E
Primary Accession	<a href="#">P11169</a>
Other Accession	<a href="#">Q8TDB8</a> , <a href="#">Q5R608</a> , <a href="#">Q9XSC2</a>
Reactivity	Human, Mouse, Rat
Predicted	Rabbit
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	53924
Antigen Region	432-463

**SLC2A3 Antibody (C-Term) - Additional Information**

**Gene ID** 6515

**Other Names**

Solute carrier family 2, facilitated glucose transporter member 3, Glucose transporter type 3, brain, GLUT-3, SLC2A3, GLUT3

**Target/Specificity**

This SLC2A3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 432-463 amino acids from human SLC2A3.

**Dilution**

WB~~1:2000  
FC~~1:25

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

SLC2A3 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**SLC2A3 Antibody (C-Term) - Protein Information**

**Name** SLC2A3 ([HGNC:11007](#))

**Function** Facilitative glucose transporter (PubMed:[26176916](#), PubMed:[32860739](#), PubMed:[9477959](#)). Can also mediate the uptake of various other monosaccharides across the cell membrane (PubMed:[26176916](#), PubMed:[9477959](#)). Mediates the uptake of glucose, 2-deoxyglucose, galactose, mannose, xylose and fucose, and probably also dehydroascorbate (PubMed:[26176916](#), PubMed:[9477959](#)). Does not mediate fructose transport (PubMed:[26176916](#), PubMed:[9477959](#)). Required for mesendoderm differentiation (By similarity).

**Cellular Location**

Cell membrane; Multi-pass membrane protein. Perikaryon {ECO:0000250|UniProtKB:Q07647}. Cell projection {ECO:0000250|UniProtKB:Q07647}. Note=Localized to densely spaced patches along neuronal processes. {ECO:0000250|UniProtKB:Q07647}

**Tissue Location**


Highly expressed in brain (PubMed:8457197). Expressed in many tissues.

**SLC2A3 Antibody (C-Term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**SLC2A3 Antibody (C-Term) - Images**



All lanes : Anti-SLC2A3 Antibody (C-Term) at 1:2000 dilution Lane 1: C6 whole cell lysate Lane 2: SH-SY5Y whole cell lysate Lane 3: human brain lysate Lane 4: mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 54 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing U-2 OS cells stained with AP22174b(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22174b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OE188374) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10, 000 events was performed.

#### **SLC2A3 Antibody (C-Term) - Background**

Facilitative glucose transporter. Probably a neuronal glucose transporter.

#### **SLC2A3 Antibody (C-Term) - References**

- Kayano T.,et al.J. Biol. Chem. 263:15245-15248(1988).
- Stuart C.A.,et al.Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases.
- Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.