

**TAF4 Antibody (C-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP22185b**

**Specification**

---

**TAF4 Antibody (C-Term) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">O00268</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>110114</b>

**TAF4 Antibody (C-Term) - Additional Information**

**Gene ID** 6874

**Other Names**

Transcription initiation factor TFIID subunit 4, RNA polymerase II TBP-associated factor subunit C, TBP-associated factor 4, Transcription initiation factor TFIID 130 kDa subunit, TAF(II)130, TAFII-130, TAFII130, Transcription initiation factor TFIID 135 kDa subunit, TAF(II)135, TAFII-135, TAFII135, TAF4, TAF2C, TAF2C1, TAF4A, TAFII130, TAFII135

**Target/Specificity**

This TAF4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1021-1062 amino acids from human TAF4.

**Dilution**

WB~~1:2000  
IHC-P~~1:25

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TAF4 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**TAF4 Antibody (C-Term) - Protein Information**

**Name** TAF4

**Synonyms** TAF2C, TAF2C1, TAF4A, TAFII130, TAFII135

**Function** The TFIID basal transcription factor complex plays a major role in the initiation of RNA polymerase II (Pol II)-dependent transcription (PubMed:[33795473](#)). TFIID recognizes and binds promoters with or without a TATA box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC) (PubMed:[33795473](#)). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and TAF13 (PubMed:[10594036](#), PubMed:[33795473](#), PubMed:[8942982](#)). TAF4 may maintain an association between the TFIID and TFIIA complexes, while bound to the promoter, together with TBP, during PIC assembly (PubMed:[33795473](#)). Potentiates transcriptional activation by the AF-2S of the retinoic acid, vitamin D3 and thyroid hormone (PubMed:[9192867](#)).

**Cellular Location**


Nucleus.

**TAF4 Antibody (C-Term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**TAF4 Antibody (C-Term) - Images**



All lanes : Anti-TAF4 Antibody (C-Term) at 1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: U-251 MG whole cell lysate Lane 3: A431 whole cell lysate Lane 4: HeLa whole cell lysate Lane 5: CCRF-CEM whole cell lysate Lane 6: HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 110 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



AP22185b staining TAF4 in human tonsil tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



AP22185b staining TAF4 in human colon tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

#### **TAF4 Antibody (C-Term) - Background**

Part of the TFIID complex, a multimeric protein complex that plays a central role in mediating promoter responses to various activators and repressors. Potentiates transcriptional activation by the AF-2S of the retinoic acid, vitamin D3 and thyroid hormone.

#### **TAF4 Antibody (C-Term) - References**

Mengus G., et al. *Genes Dev.* 11:1381-1395(1997).  
Deloukas P., et al. *Nature* 414:865-871(2001).

Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Tanese N.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:13611-13616(1996).  
Damania B.,et al.Genes Dev. 10:1369-1381(1996).