

TPD52 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22193a

Specification

TPD52 Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	P55327
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	24327

TPD52 Antibody (N-Term) - Additional Information

Gene ID 7163

Other Names

Tumor protein D52, Protein N8, TPD52

Target/Specificity

This TPD52 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 11-44 amino acids from human TPD52.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TPD52 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

TPD52 Antibody (N-Term) - Protein Information

Name TPD52

Tissue Location

Isoform 2 is expressed in colon, breast, prostate, pancreas and kidney tumor cell lines. Isoform 2 is expressed at high levels in kidney, prostate, brain, small intestine and pancreas, at moderate levels in placenta and colon, at low levels in lung, liver and heart, and at very low levels in spleen,


thymus, peripheral mononuclear blood cells, testis and ovary.

TPD52 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TPD52 Antibody (N-Term) - Images



All lanes : Anti-TPD52 Antibody (N-Term) at 1:2000 dilution Lane 1: mouse brain lysate Lane 2: mouse kidney lysate Lane 3: THP-1 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

TPD52 Antibody (N-Term) - References

- Byrne J.A.,et al.Cancer Res. 55:2896-2903(1995).
Chen S.-L.,et al.Oncogene 12:741-751(1996).
Wang R.,et al.Cancer Res. 64:1589-1594(2004).
Wang R.,et al.Biochem. Biophys. Res. Commun. 389:455-460(2009).
Ota T.,et al.Nat. Genet. 36:40-45(2004).