

OVGP1 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22205a**Specification**

OVGP1 Antibody (N-Term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	O12889
Other Accession	P36718
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	75421

OVGP1 Antibody (N-Term) - Additional Information**Gene ID** 5016**Other Names**

Oviduct-specific glycoprotein, Estrogen-dependent oviduct protein, Mucin-9, Oviductal glycoprotein, Oviductin, OVGP1, MUC9, OGP

Target/Specificity

This OVGP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 107-137 amino acids from human OVGP1.

Dilution

WB~~1:2000

IHC-P~~1:25

FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OVGP1 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

OVGP1 Antibody (N-Term) - Protein Information**Name** OVGP1

Synonyms MUC9, OGP

Function Binds to oocyte zona pellucida in vivo. May play a role in the fertilization process and/or early embryonic development.

Cellular Location

Cytoplasmic vesicle, secretory vesicle. Note=Secretory granules

Tissue Location


Oviduct.

OVGP1 Antibody (N-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


OVGP1 Antibody (N-Term) - Images



All lanes : Anti-OVGP1 Antibody (N-Term) at 1:2000 dilution Lane 1: human placenta lysate Lane 2: human uterus lysate Lane 3: human ovary lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 75 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



AP22205a staining OVGP1 in human fallopian tube tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing Hela cells stained with AP22205a (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22205a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed (1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

OVGP1 Antibody (N-Term) - Background

Binds to oocyte zona pellucida in vivo. May play a role in the fertilization process and/or early embryonic development.

OVGP1 Antibody (N-Term) - References

Arias E.B., et al. *Biol. Reprod.* 51:685-694(1994).
Jaffe R.C., et al. Submitted (JUL-1996) to the EMBL/GenBank/DDBJ databases.

Gregory S.G.,et al.Nature 441:315-321(2006).
Sjoebloom T.,et al.Science 314:268-274(2006).