

RPS2 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22208a

Specification

RPS2 Antibody (N-Term) - Product Information

| | |
|-------------------|--|
| Application | WB, FC,E |
| Primary Accession | P15880 |
| Other Accession | O18789 , P46791 , Q90YS3 , P25444 , P27952 , G1SWM1 |
| Reactivity | Human, Mouse |
| Predicted | Bovine, Hamster, Rabbit, Rat |
| Host | Rabbit |
| Clonality | polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 31324 |

RPS2 Antibody (N-Term) - Additional Information

Gene ID 6187

Other Names

40S ribosomal protein S2, 40S ribosomal protein S4, Protein LLRep3, RPS2, RPS4

Target/Specificity

This RPS2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 45-79 amino acids from human RPS2.

Dilution

WB~~1:2000
FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RPS2 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

RPS2 Antibody (N-Term) - Protein Information

Name RPS2

Synonyms RPS4

Function Component of the ribosome, a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed:[23636399](#)). The small ribosomal subunit (SSU) binds messenger RNAs (mRNAs) and translates the encoded message by selecting cognate aminoacyl-transfer RNA (tRNA) molecules (PubMed:[23636399](#)). The large subunit (LSU) contains the ribosomal catalytic site termed the peptidyl transferase center (PTC), which catalyzes the formation of peptide bonds, thereby polymerizing the amino acids delivered by tRNAs into a polypeptide chain (PubMed:[23636399](#)). The nascent polypeptides leave the ribosome through a tunnel in the LSU and interact with protein factors that function in enzymatic processing, targeting, and the membrane insertion of nascent chains at the exit of the ribosomal tunnel (PubMed:[23636399](#)). Plays a role in the assembly and function of the 40S ribosomal subunit (By similarity). Mutations in this protein affects the control of translational fidelity (By similarity). Involved in nucleolar processing of pre-18S ribosomal RNA and ribosome assembly (By similarity).

Cellular Location


Cytoplasm. Nucleus, nucleolus. Note=Probably localized to nucleolus and cytoplasm in complex with ZNF277.

RPS2 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RPS2 Antibody (N-Term) - Images



All lanes : Anti-RPS2 Antibody (N-Term) at 1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: NIH/3T3 whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated

at 1/10000 dilution. Predicted band size : 31 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing Jurkat cells stained with AP22208a(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22208a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

RPS2 Antibody (N-Term) - References

- Slynn G.,et al.Nucleic Acids Res. 18:681-681(1990).
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
- Vladimirov S.N.,et al.Eur. J. Biochem. 239:144-149(1996).
- Swiercz R.,et al.Biochem. J. 386:85-91(2005).