

**GGT3P Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP22260c****Specification**

---

**GGT3P Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">A6NGU5</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	61502
Antigen Region	176-207

**GGT3P Antibody (Center) - Additional Information****Target/Specificity**

This GGT3P antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 176-207 amino acids from the Central region of human GGT3P.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GGT3P Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**GGT3P Antibody (Center) - Protein Information**

**Name** GGT3P ([HGNC:4252](#))

**Synonyms** GGT3

**Function** Hydrolyzes and transfers gamma-glutamyl moieties from glutathione and other gamma-glutamyl compounds to acceptors.

**Cellular Location**


Membrane {ECO:0000250|UniProtKB:P19440}; Single-pass type II membrane protein {ECO:0000250|UniProtKB:P07314}

## GGT3P Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## GGT3P Antibody (Center) - Images



Anti-GGT3P Antibody (Center) at 1:2000 dilution + Human kidney lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 62 kDa Blocking/Dilution buffer: 5% NFDN/TBST.