

IL11 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22261a

Specification

IL11 Antibody (N-Term) - Product Information

Application	WB, FC,E
Primary Accession	P20809
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	21429

IL11 Antibody (N-Term) - Additional Information

Gene ID 3589

Other Names

Interleukin-11, IL-11, Adipogenesis inhibitory factor, AGIF, Oprelvekin, IL11

Target/Specificity

This IL11 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 27-61 amino acids from human IL11.

Dilution

WB~~1:2000

FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IL11 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

IL11 Antibody (N-Term) - Protein Information

Name IL11 ([HGNC:5966](#))

Function Cytokine that stimulates the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells and induces megakaryocyte maturation resulting in increased platelet production (PubMed:[2145578](#)). Also promotes the proliferation of hepatocytes in response

to liver damage. Binding to its receptor formed by IL6ST and IL11RA activates a signaling cascade that promotes cell proliferation (PubMed:[12919066](#)). Signaling leads to the activation of intracellular protein kinases and the phosphorylation of STAT3. The interaction with the membrane-bound IL11RA and IL6ST stimulates 'classic signaling', whereas the binding of IL11 and soluble IL11RA to IL6ST stimulates 'trans-signaling' (PubMed:[30279168](#)).

Cellular Location


Secreted

IL11 Antibody (N-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IL11 Antibody (N-Term) - Images



Anti-IL11 Antibody (N-Term) at 1:2000 dilution + DU145 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 21 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



Overlay histogram showing PC-3 cells stained with AP22261a (green line). The cells were fixed with 2% paraformaldehyde and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at Room temperature. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

IL11 Antibody (N-Term) - Background

Cytokine that stimulates the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells and induces megakaryocyte maturation resulting in increased platelet production (PubMed:2145578). Also promotes the proliferation of hepatocytes in response to liver damage. Binding to its receptor formed by IL6ST and either IL11RA1 or IL11RA2 activates a signaling cascade that promotes cell proliferation (PubMed:12919066). Signaling leads to the activation of intracellular protein kinases and the phosphorylation of STAT3.

IL11 Antibody (N-Term) - References

- Paul S.R., et al. Proc. Natl. Acad. Sci. U.S.A. 87:7512-7516(1990).
- Kawashima I., et al. FEBS Lett. 283:199-202(1991).
- McKinley D., et al. Genomics 13:814-819(1992).
- Ota T., et al. Nat. Genet. 36:40-45(2004).
- Grimwood J., et al. Nature 428:529-535(2004).