

Wee1(S123) Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22278a**Specification**

Wee1(S123) Antibody - Product Information

Application	WB,E
Primary Accession	P30291
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	71597

Wee1(S123) Antibody - Additional Information**Gene ID** 7465**Other Names**

Wee1-like protein kinase, WEE1hu, 2.7.10.2, Wee1A kinase, WEE1

Target/Specificity

This Wee1(S123) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 96-130 amino acids from human Wee1.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Wee1(S123) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Wee1(S123) Antibody - Protein Information**Name** WEE1 {ECO:0000303|PubMed:8348613, ECO:0000312|HGNC:HGNC:12761}**Function** Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15' (PubMed:[15070733](#), PubMed:[7743995](#), PubMed:[8348613](#), PubMed:[8428596](#)). Specifically phosphorylates and inactivates cyclin

B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase (PubMed:[7743995](#), PubMed:[8348613](#), PubMed:[8428596](#)). Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and phosphorylation of monomeric CDK1 does not occur (PubMed:[7743995](#), PubMed:[8348613](#), PubMed:[8428596](#)). Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated (PubMed:[7743995](#)). A correlated decrease in protein level occurs at M/G1 phase, probably due to its degradation (PubMed:[7743995](#)).

Cellular Location


Nucleus.

Wee1(S123) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Wee1(S123) Antibody - Images



All lanes : Anti-Wee1(S123) Antibody at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 72 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

Wee1(S123) Antibody - Background

Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15'. Specifically phosphorylates and inactivates cyclin B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase.

Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and phosphorylation of monomeric CDK1 does not occur. Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated. A correlated decrease in protein level occurs at M/G1 phase, probably due to its degradation.

Wee1(S123) Antibody - References

- Watanabe N., et al. EMBO J. 14:1878-1891(1995).
Cichutek A., et al. Cytogenet. Cell Genet. 93:277-283(2001).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Taylor T.D., et al. Nature 440:497-500(2006).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.