

CDKN2A antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22403a

Specification

CDKN2A antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB,E |
| Primary Accession | P42771 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | polyclonal |
| Isotype | Rabbit Ig |
| Calculated MW | 16533 |

CDKN2A antibody - Additional Information

Gene ID 1029

Other Names

Cyclin-dependent kinase inhibitor 2A {ECO:0000312|HGNC:HGNC:1787}, Cyclin-dependent kinase 4 inhibitor A, CDK4I, Multiple tumor suppressor 1, MTS-1, p16-INK4a, p16-INK4, p16INK4A, CDKN2A (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=1787), CDKN2, MTS1

Target/Specificity

This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDKN2A antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CDKN2A antibody - Protein Information

Name CDKN2A ([HGNC:1787](#))

Synonyms CDKN2, MTS1

Function Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.

Cellular Location

Cytoplasm. Nucleus

Tissue Location


Widely expressed but not detected in brain or skeletal muscle. Isoform 3 is pancreas-specific

CDKN2A antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDKN2A antibody - Images



All lanes: Anti-CDKN2A antibody at 1:2000 dilution Lane 1: 293 whole cell lysate Lane 2: 293T whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 17 KDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes: Anti-CDKN2A antibody at 1:1000 dilution Lane 1: 293 whole cell lysate Lane 2: 293T whole cell lysate Lane 3: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 17 KDa Blocking/Dilution buffer: 5% NFD/MTBST.

CDKN2A antibody - Background

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.

CDKN2A antibody - References

Serrano M., et al. Nature 366:704-707(1993).
Robertson K.D., et al. Oncogene 18:3810-3820(1999).
Kitagawa Y., et al. J. Biol. Chem. 277:46289-46297(2002).
Lin Y.C., et al. Oncogene 26:7017-7027(2007).
Humphray S.J., et al. Nature 429:369-374(2004).