

**FGL1 antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP22409a**

**Specification**

---

**FGL1 antibody - Product Information**

Application	<b>WB,E</b>
Primary Accession	<a href="#">O08830</a>
Host	<b>Rabbit</b>
Clonality	<b>polyclonal</b>
Isotype	<b>Rabbit Ig</b>
Calculated MW	<b>36379</b>

**FGL1 antibody - Additional Information**

**Gene ID** 2267

**Other Names**

Fibrinogen-like protein 1, HP-041, Hepassocin, HPS, Hepatocyte-derived fibrinogen-related protein 1, HFREP-1, Liver fibrinogen-related protein 1 {ECO:0000303|Ref.3}, LFIRE-1 {ECO:0000303|Ref.3}, FGL1 {ECO:0000303|PubMed:18039467, ECO:0000312|HGNC:HGNC:3695}

**Target/Specificity**

This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

FGL1 antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**FGL1 antibody - Protein Information**

**Name** FGL1 {ECO:0000303|PubMed:18039467, ECO:0000312|HGNC:HGNC:3695}

**Function** Immune suppressive molecule that inhibits antigen-specific T- cell activation by acting as a major ligand of LAG3 (PubMed:[30580966](#)). Responsible for LAG3 T-cell inhibitory function (PubMed:[30580966](#)). Binds LAG3 independently from MHC class II (MHC-II) (PubMed:[30580966](#)). Secreted by, and promotes growth of, hepatocytes (PubMed:[11470158](#), PubMed:[19880967](#)).

**Cellular Location**

Secreted. Note=Secreted in the blood plasma


**Tissue Location**

Under normal conditions, liver-specific.

**FGL1 antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**FGL1 antibody - Images**

All lanes: Anti-FGL1 antibody at 1:2000 dilution + HepG2 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 32 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

**FGL1 antibody - Background**

Immune suppressive molecule that inhibits antigen-specific T- cell activation by acting as a major ligand of LAG3 (PubMed:30580966). Responsible for LAG3 T-cell inhibitory function (PubMed:30580966). Binds LAG3 independently from MHC class II (MHC-II) (PubMed:30580966). Secreted by, and promotes growth of, hepatocytes (PubMed:11470158, PubMed:19880967).

**FGL1 antibody - References**

Yamamoto T.,et al.Biochem. Biophys. Res. Commun. 193:681-687(1993).  
Hara H.,et al.Biochim. Biophys. Acta 1520:45-53(2001).  
Yan J.,et al.Submitted (JUL-1999) to the EMBL/GenBank/DBJ databases.

Kalnine N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.  
Ota T., et al. Nat. Genet. 36:40-45(2004).