

CD38
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22413a

Specification

CD38 - Product Information

Application	IHC-P,E
Primary Accession	P28907
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	34328

CD38 - Additional Information

Gene ID 952

Other Names

ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, 3.2.2.-, 3.2.2.6, 2'-phospho-ADP-ribosyl cyclase, 2'-phospho-ADP-ribosyl cyclase/2'-phospho-cyclic-ADP-ribose transferase, 2.4.99.20, 2'-phospho-cyclic-ADP-ribose transferase, ADP-ribosyl cyclase 1, ADPRC 1, Cyclic ADP-ribose hydrolase 1, cADPR hydrolase 1, T10, CD38, CD38

Target/Specificity

This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.

Dilution

IHC-P~~1ug/ml

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CD38 is for research use only and not for use in diagnostic or therapeutic procedures.

CD38 - Protein Information

Name CD38

Function Synthesizes cyclic ADP-ribose (cADPR), a second messenger for glucose-induced insulin secretion (PubMed:[7961800](#), PubMed:[8253715](#)). Synthesizes the Ca(2+) mobilizer

nicotinate-adenine dinucleotide phosphate, NAADP(+), from 2'-phospho-cADPR and nicotinic acid, as well as from NADP(+) and nicotinic acid. At both pH 5.0 and pH 7.4 preferentially transforms 2'-phospho-cADPR into NAADP(+), while preferentially cleaving NADP(+) to cADPR and ADPRP rather than into NADDP(+) (PubMed:[16690024](#)). Has cADPR hydrolase activity (PubMed:[7961800](#), PubMed:[8253715](#)).

Cellular Location

Cell surface. Membrane; Single-pass type II membrane protein

Tissue Location


Expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma.

CD38 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD38 - Images



Immunohistochemical analysis of paraffin-embedded Human Tonsil tissue using CD38 antibody (C-term) performed on the Abcarta® FAIP-48T Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9.0). Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:ADR005] was used as the secondary antibody.

CD38 - Background

Synthesizes cyclic ADP-ribose (cADPR), a second messenger for glucose-induced insulin secretion (PubMed:8253715, PubMed:7961800). Synthesizes the Ca²⁺ mobilizer nicotinate-adenine dinucleotide phosphate, NAADP(+), from 2'-phospho-cADPR and nicotinic acid, as well as from NADP(+) and nicotinic acid. At both pH 5.0 and pH 7.4 preferentially transforms 2'-phospho-cADPR into NAADP(+), while preferentially cleaving NADP(+) to cADPR and ADPRP rather than into

NADDP(+) (PubMed:16690024). Has cADPR hydrolase activity (PubMed:8253715, PubMed:7961800).

CD38 - References

- Jackson D.G.,et al.J. Immunol. 144:2811-2815(1990).
Nata K.,et al.Gene 186:285-292(1997).
States D.J.,et al.Trends Biochem. Sci. 17:495-495(1992).
Takasawa S.,et al.J. Biol. Chem. 268:26052-26054(1993).
Tohgo A.,et al.J. Biol. Chem. 269:28555-28557(1994).