

Neurofilament-L antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22455a

Specification

Neurofilament-L antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB,E |
| Primary Accession | P07196 |
| Reactivity | Human, Mouse |
| Host | Rabbit |
| Clonality | polyclonal |
| Isotype | Rabbit Ig |
| Calculated MW | 61517 |

Neurofilament-L antibody - Additional Information

Gene ID 4747

Other Names

Neurofilament light polypeptide, NF-L, 68 kDa neurofilament protein, Neurofilament triplet L protein, NEFL, NF68, NFL

Target/Specificity

This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Neurofilament-L antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Neurofilament-L antibody - Protein Information

Name NEFL

Synonyms NF68, NFL

Function Neurofilaments usually contain three intermediate filament proteins: NEFL, NEFM, and

NEFH which are involved in the maintenance of neuronal caliber. May additionally cooperate with the neuronal intermediate filament proteins PRPH and INA to form neuronal filamentous networks (By similarity).

Cellular Location


Cell projection, axon {ECO:0000250|UniProtKB:P08551}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P08551}

Neurofilament-L antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Neurofilament-L antibody - Images



All lanes: Anti-Neurofilament-L antibody at 1:2000 dilution Lane 1: Human brain lysate Lane 2: Mouse brain lysate Lane 3: Rat brain lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 62 KDa Blocking/Dilution buffer: 5% NFD/MTBST.

Neurofilament-L antibody - Background

Neurofilaments usually contain three intermediate filament proteins: NEFL, NEFM, and NEFH which are involved in the maintenance of neuronal caliber. May additionally cooperate with the neuronal intermediate filament proteins PRPH and INA to form neuronal filamentous networks (By similarity).

Neurofilament-L antibody - References

Julien J.-P., et al. Biochim. Biophys. Acta 909:10-20(1987).

Perez-Olle R., et al. J. Cell Sci. 115:4937-4946(2002).
Nusbaum C., et al. Nature 439:331-335(2006).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Pospelov V.A., et al. Cell Growth Differ. 5:187-196(1994).