

**LAG3 antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP22459a****Specification**

---

**LAG3 antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P18627</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	57449

**LAG3 antibody (N-term) - Additional Information****Gene ID** 3902**Other Names**

Lymphocyte activation gene 3 protein, LAG-3, CD223, Secreted lymphocyte activation gene 3 protein, sLAG-3, LAG3 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=6476" target="\_blank">HGNC:6476</a>), FDC

**Target/Specificity**

This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.

**Dilution**

WB~~1:1000  
E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

LAG3 antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**LAG3 antibody (N-term) - Protein Information****Name** LAG3 ([HGNC:6476](#))

## Synonyms FDC

**Function** Lymphocyte activation gene 3 protein: Inhibitory receptor on antigen activated T-cells (PubMed:[20421648](#), PubMed:[7805750](#), PubMed:[8647185](#)). Delivers inhibitory signals upon binding to ligands, such as FGL1 (By similarity). FGL1 constitutes a major ligand of LAG3 and is responsible for LAG3 T-cell inhibitory function (By similarity). Following TCR engagement, LAG3 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1, possibly by acting as a coreceptor for PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed:[20421648](#), PubMed:[7805750](#), PubMed:[8647185](#)). Also mediates immune tolerance: constitutively expressed on a subset of regulatory T-cells (Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity). Binds MHC class II (MHC-II); the precise role of MHC-II-binding is however unclear (PubMed:[8647185](#)).

## Cellular Location

[Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein

## Tissue Location


Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.

## LAG3 antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## LAG3 antibody (N-term) - Images



All lanes: Anti-LAG3 antibody (N-term) at 1:1000 dilution Lane 1: A2058 whole cell lysate Lane 2: K562 whole cell lysate Lane 3: Rat thymus lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 57 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

### **LAG3 antibody (N-term) - Background**

Lymphocyte activation gene 3 protein: Inhibitory receptor on antigen activated T-cells (PubMed:20421648, PubMed:7805750, PubMed:8647185). Delivers inhibitory signals upon binding to ligands, such as FGL1 (By similarity). FGL1 constitutes a major ligand of LAG3 and is responsible for LAG3 T-cell inhibitory function (By similarity). Following TCR engagement, LAG3 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1, possibly by acting as a coreceptor for PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed:20421648, PubMed:7805750, PubMed:8647185). Also mediates immune tolerance: constitutively expressed on a subset of regulatory T-cells (Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity). Binds MHC class II (MHC-II); the precise role of MHC-II-binding is however unclear (PubMed:8647185).

### **LAG3 antibody (N-term) - References**

Triebel F.,et al.J. Exp. Med. 171:1393-1405(1990).  
Triebel F.,et al.Submitted (SEP-2001) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Scherer S.E.,et al.Nature 440:346-351(2006).  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.