

## **ZIC4 Antibody (C-term)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2722b

### **Specification**

## **ZIC4 Antibody (C-term) - Product Information**

**Application** WB,E **Primary Accession Q8N9L1** Reactivity Human **Rabbit** Host Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 36567 Antigen Region 299-334

### ZIC4 Antibody (C-term) - Additional Information

#### **Gene ID 84107**

### **Other Names**

Zinc finger protein ZIC 4, Zinc finger protein of the cerebellum 4, ZIC4

# **Target/Specificity**

This ZIC4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 299-334 amino acids from the C-terminal region of human ZIC4.

#### **Dilution**

WB~~1:1000

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

## **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

ZIC4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### **ZIC4 Antibody (C-term) - Protein Information**

Name ZIC4

Function Binds to DNA.

**Cellular Location** 




Nucleus.

## **ZIC4 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## ZIC4 Antibody (C-term) - Images



Western blot analysis of ZIC4 (arrow) using rabbit polyclonal ZIC4 Antibody (C-term) (Cat.#AP2722b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the ZIC4 gene (Lane 2) (Origene Technologies).

## ZIC4 Antibody (C-term) - Background

ZIC4 is a member of the ZIC family of C2H2-type zinc finger proteins. Members of this family are important during development, and have been associated with X-linked visceral heterotaxy and holoprosencephaly type 5.

## **ZIC4 Antibody (C-term) - References**

Ishiguro, A., Biochem. Biophys. Res. Commun. 324 (1), 302-307 (2004) Grinberg, I., Nat. Genet. 36 (10), 1053-1055 (2004) Aruga, J., Gene 172 (2), 291-294 (1996)