

**AKR1A1 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2734b****Specification**

---

**AKR1A1 Antibody (C-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">P14550</a>
Other Accession	<a href="#">P51635</a> , <a href="#">P50578</a> , <a href="#">Q9JII6</a> , <a href="#">Q3ZCJ2</a> , <a href="#">Q6IAZ4</a>
Reactivity	Human
Predicted	Bovine, Mouse, Pig, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36573
Antigen Region	293-325

**AKR1A1 Antibody (C-term) - Additional Information****Gene ID** 10327**Other Names**

Alcohol dehydrogenase [NADP(+)], Aldehyde reductase, Aldo-keto reductase family 1 member A1, AKR1A1, ALDR1, ALR

**Target/Specificity**

This AKR1A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 293-325 amino acids from the C-terminal region of human AKR1A1.

**Dilution**WB~~1:1000  
IHC-P~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

AKR1A1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**AKR1A1 Antibody (C-term) - Protein Information****Name** AKR1A1

## Synonyms ALDR1, ALR

**Function** Catalyzes the NADPH-dependent reduction of a wide variety of carbonyl-containing compounds to their corresponding alcohols (PubMed:[10510318](#), PubMed:[30538128](#)). Displays enzymatic activity towards endogenous metabolites such as aromatic and aliphatic aldehydes, ketones, monosaccharides and bile acids, with a preference for negatively charged substrates, such as glucuronate and succinic semialdehyde (PubMed:[10510318](#), PubMed:[30538128](#)). Functions as a detoxifying enzyme by reducing a range of toxic aldehydes (By similarity). Reduces methylglyoxal and 3-deoxyglucosone, which are present at elevated levels under hyperglycemic conditions and are cytotoxic (By similarity). Involved also in the detoxification of lipid-derived aldehydes like acrolein (By similarity). Plays a role in the activation of procarcinogens, such as polycyclic aromatic hydrocarbon trans-dihydrodiols, and in the metabolism of various xenobiotics and drugs, including the anthracyclines doxorubicin (DOX) and daunorubicin (DAUN) (PubMed:[11306097](#), PubMed:[18276838](#)). Also acts as an inhibitor of protein S-nitrosylation by mediating degradation of S-nitroso-coenzyme A (S-nitroso-CoA), a cofactor required to S-nitrosylate proteins (PubMed:[30538128](#)). S-nitroso-CoA reductase activity is involved in reprogramming intermediary metabolism in renal proximal tubules, notably by inhibiting protein S-nitrosylation of isoform 2 of PKM (PKM2) (By similarity). Also acts as a S-nitroso- glutathione reductase by catalyzing the NADPH-dependent reduction of S- nitrosoglutathione (PubMed:[31649033](#)). Displays no reductase activity towards retinoids (By similarity).

## Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9JII6}. Apical cell membrane {ECO:0000250|UniProtKB:Q9JII6}

## Tissue Location


Widely expressed. Highly expressed in kidney, salivary gland and liver. Detected in trachea, stomach, brain, lung, prostate, placenta, mammary gland, small intestine and lung


## AKR1A1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## AKR1A1 Antibody (C-term) - Images






Western blot analysis of anti-AKR1A1 Antibody (C-term) (Cat.#AP2734b) in Y79 cell line lysates (35ug/lane). AKR1A1 (arrow) was detected using the purified Pab.



Western blot analysis of AKR1A1 (arrow) using rabbit polyclonal AKR1A1 Antibody (C-term) (Cat.#AP2734b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the AKR1A1 gene.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with AKR1A1 antibody (C-term) (Cat.#AP2734b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

**AKR1A1 Antibody (C-term) - Background**

AKR1A1 is a member of the aldo/keto reductase superfamily, which consists of more than 40

known enzymes and proteins. This member, also known as aldehyde reductase, is involved in the reduction of biogenic and xenobiotic aldehydes and is present in virtually every tissue.

#### **AKR1A1 Antibody (C-term) - References**

- Steuber, H.J. Mol. Biol. 379 (5), 991-1016 (2008)  
Bohren, K.M., Biochim. Biophys. Acta 1748 (2), 201-212 (2005)  
El-Kabbani, O., Acta Crystallogr. D Biol. Crystallogr. 50 (PT 6), 859-868 (1994)