

**ALDOC Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2736b****Specification**

---

**ALDOC Antibody (C-term) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">P09972</a>
Other Accession	<a href="#">P09117</a> , <a href="#">P05063</a> , <a href="#">Q9GKW3</a>
Reactivity	<b>Human</b>
Predicted	<b>Monkey, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>39456</b>
Antigen Region	<b>300-333</b>

**ALDOC Antibody (C-term) - Additional Information****Gene ID** 230**Other Names**

Fructose-bisphosphate aldolase C, Brain-type aldolase, ALDOC, ALDC

**Target/Specificity**

This ALDOC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 300-333 amino acids from the C-terminal region of human ALDOC.

**Dilution**WB~~1:1000  
IHC-P~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ALDOC Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**ALDOC Antibody (C-term) - Protein Information****Name** ALDOC


## Synonyms ALDC

### ALDOC Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### ALDOC Antibody (C-term) - Images



Western blot analysis of lysates from human brain, mouse cerebellum and rat brain tissue lysate (from left to right), using ALDOC Antibody (C-term)(Cat. #AP2736b). AP2736b was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



Formalin-fixed and paraffin-embedded human Brain, Cerebellum tissue reacted with ALDOC antibody (C-term) (Cat.#AP2736b), which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

#### **ALDOC Antibody (C-term) - Background**

ALDOC is a member of the fructose-biphosphate aldolase protein family. Expressed specifically in the hippocampus and Purkinje cells of the brain, this protein is a glycolytic enzyme that catalyzes the reversible aldol cleavage of fructose-1,6-biphosphate and fructose 1-phosphate to dihydroxyacetone phosphate and either glyceraldehyde-3-phosphate or glyceraldehyde, respectively.

#### **ALDOC Antibody (C-term) - References**

- Buono,P.,FEBS Lett. 578 (3), 337-344 (2004)
- Buono,P.,Nucleic Acids Res. 16 (10), 4733 (1988)
- Rottmann,W.H.,Proc. Natl. Acad. Sci. U.S.A. 56 (4), 1275-1282 (1966)