

Vimentin Antibody (S82)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2739a**Specification**

Vimentin Antibody (S82) - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	P08670
Other Accession	P31000 , P20152 , O4R4X4 , P48670 , P48616 , O9MZA9
Reactivity	Human, Mouse
Predicted	Bovine, Hamster, Monkey, Rat, Sheep
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	63-90

Vimentin Antibody (S82) - Additional Information**Gene ID** 7431**Other Names**
Vimentin, VIM**Target/Specificity**

This Vimentin antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 63-90 amino acids from human Vimentin.

DilutionIF~~1:25
WB~~1:1000
IHC-P~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Vimentin Antibody (S82) is for research use only and not for use in diagnostic or therapeutic procedures.

Vimentin Antibody (S82) - Protein Information**Name** VIM ([HGNC:12692](#))

Function Vimentins are class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. Vimentin is attached to the nucleus, endoplasmic reticulum, and mitochondria, either laterally or terminally. Plays a role in cell directional movement, orientation, cell sheet organization and Golgi complex polarization at the cell migration front (By similarity). Protects SCRIB from proteasomal degradation and facilitates its localization to intermediate filaments in a cell contact-mediated manner (By similarity).

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Nucleus matrix {ECO:0000250|UniProtKB:P31000}. Cell membrane {ECO:0000250|UniProtKB:P20152}

Tissue Location


Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone-independent mammary carcinoma cell lines.

Vimentin Antibody (S82) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Vimentin Antibody (S82) - Images




Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized U-2 OS (human osteosarcoma cell line) cells labeling Vimentin with AP2739a at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (1583138) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm and weak nucleus staining on U-2 OS cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red). The nuclear counter stain is DAPI (blue).



Western blot analysis of Vimentin-S82 (Cat. #AP2739a) in HeLa cell line lysates (35ug/lane). Vimentin (arrow) was detected using the purified Pab.




Western blot analysis of VIM (arrow) using rabbit polyclonal Vimentin Antibody (S82) (Cat.#AP2739a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the VIM gene (Lane 2) (Origene Technologies).



All lanes : Anti-Vimentin Antibody (S82) at 1:1000 dilution Lane 1: 293 whole cell lysate Lane 2:

Hela whole cell lysate Lane 3: U-2OS whole cell lysate Lane 4: C2C12 whole cell lysate Lane 5: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 54 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human breast carcinoma tissue reacted with Vimentin Antibody (S82) (Cat.#AP2739a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Vimentin Antibody (S82) - Background

Along with the microfilaments (actins) and microtubules (tubulins), the intermediate filaments represent a third class of well-characterized cytoskeletal elements. The subunits display a tissue-specific pattern of expression. Desmin (MIM 125660) is the subunit specific for muscle and vimentin the subunit specific for mesenchymal tissue.

Vimentin Antibody (S82) - References

References for protein:

1. Whipple, R.A., Cancer Res. 68 (14), 5678-5688 (2008)
2. Garcia-Verdugo, I., Biochemistry 47 (18), 5127-5138 (2008)
3. Merdes, A., J. Cell Biol. 115 (2), 397-410 (1991)

References for SY5Y (SH-SY5Y; ATCC#CRL-2266): 1. Ross RA, et al. Coordinate morphological and biochemical interconversion of human neuroblastoma cells. J. Natl. Cancer Inst. 71: 741-749, 1983. [PubMed: 6137586]; 2. Biedler JL, et al. Multiple neurotransmitter synthesis by human neuroblastoma cell lines and clones. Cancer Res. 38: 3751-3757, 1978. [PubMed: 29704]

Vimentin Antibody (S82) - Citations

- [Isolation and feeder-free primary culture of four cell types from a single human skin sample](#)
- [Pirfenidone inhibits epithelial-mesenchymal transition in keloid keratinocytes](#)