

FUT4 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2773c**Specification**

FUT4 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P22083
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	59084
Antigen Region	242-271

FUT4 Antibody (Center) - Additional Information**Gene ID** 2526**Other Names**

Alpha-(1, 3)-fucosyltransferase 4, 241-, ELAM-1 ligand fucosyltransferase, Fucosyltransferase 4, Fucosyltransferase IV, Fuc-TIV, FucT-IV, Galactoside 3-L-fucosyltransferase, FUT4, ELFT, FCT3A

Target/Specificity

This FUT4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 242-271 amino acids from the Central region of human FUT4.

DilutionWB~~1:1000
IHC-P~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FUT4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

FUT4 Antibody (Center) - Protein Information**Name** FUT4 {ECO:0000303|PubMed:29593094}**Function** [Isoform Short]: Catalyzes alpha(1->3) linkage of fucosyl moiety transferred from

GDP-beta-L-fucose to N-acetyl glucosamine (GlcNAc) within type 2 lactosamine (LacNAc, Gal-beta(1->4)GlcNAc) glycan attached to N- or O-linked glycoproteins (PubMed:[1702034](#), PubMed:[1716630](#), PubMed:[29593094](#)). Robustly fucosylates nonsialylated distal LacNAc unit of the poly lactosamine chain to form Lewis X antigen (CD15), a glycan determinant known to mediate important cellular functions in development and immunity. Fucosylates with lower efficiency sialylated LacNAc acceptors to form sialyl Lewis X and 6- sulfo sialyl Lewis X determinants that serve as recognition epitopes for C-type lectins (PubMed:[1716630](#), PubMed:[29593094](#)). Together with FUT7 contributes to SELE, SELL and SELP selectin ligand biosynthesis and selectin-dependent lymphocyte homing, leukocyte migration and blood leukocyte homeostasis (By similarity). In a cell type specific manner, may also fucosylate the internal LacNAc unit of the poly lactosamine chain to form VIM-2 antigen that serves as recognition epitope for SELE (PubMed:[11278338](#), PubMed:[1716630](#)).

Cellular Location

Golgi apparatus, Golgi stack membrane; Single- pass type II membrane protein.
Note=Membrane-bound form in trans cisternae of Golgi

Tissue Location


[Isoform Short]: Expressed at low levels in bone marrow-derived mesenchymal stem cells.

FUT4 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FUT4 Antibody (Center) - Images



FUT4 Antibody (Center) (Cat.#AP2773c) western blot analysis in ZR-75-1 cell line lysates (35ug/lane). This demonstrates the FUT4 antibody detected the FUT4 protein (arrow).



FUT4 Antibody (Center) (Cat. #AP2773c) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of FUT4 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

FUT4 Antibody (Center) - Background

FUT4 catalyzes the synthesis of the non-sialylated antigen, Lewis x (CD15). It has been detected in human embryos (5-10 weeks) suggesting a role in development.

FUT4 Antibody (Center) - References

Zhang, Z., *Biochim. Biophys. Acta* 1783 (2), 287-296 (2008)
Cruse, J.M., *Exp. Mol. Pathol.* 83 (2), 274-276 (2007)