

APRT Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2893a

Specification

APRT Antibody (N-term) - Product Information

Application	IF, WB, IHC-P, FC,E
Primary Accession	P07741
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	12-40

APRT Antibody (N-term) - Additional Information

Gene ID 353

Other Names

Adenine phosphoribosyltransferase, APRT, APRT

Target/Specificity

This APRT antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 12-40 amino acids from the N-terminal region of human APRT.

Dilution

IF~~1:10~50
WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

APRT Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

APRT Antibody (N-term) - Protein Information

Name APRT ([HGNC:626](#))

Function Catalyzes a salvage reaction resulting in the formation of AMP, that is energetically less

costly than de novo synthesis.

Cellular Location


Cytoplasm.

APRT Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


APRT Antibody (N-term) - Images




Fluorescent image of A549 cell stained with APRT Antibody (N-term)(Cat#AP2893a/SA091009AF).A549 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with APRT primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C).Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C).APRT immunoreactivity is localized to Cytoplasm and Nucleus significantly.



APRT Antibody (N-term) (Cat. #AP2893a) western blot analysis in A2058 cell line lysates (35ug/lane). This demonstrates the APRT antibody detected the APRT protein (arrow).



Formalin-fixed and paraffin-embedded human hepatocarcinoma with APRT Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of HepG2 cells using APRT Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

APRT Antibody (N-term) - Background

A conserved feature of APRT is the distribution of CpG dinucleotides. This enzyme catalyzes the formation of AMP and inorganic pyrophosphate from adenine and 5-phosphoribosyl-1-pyrophosphate (PRPP). It also produces adenine as a by-product of the polyamine biosynthesis pathway. A homozygous deficiency in this enzyme causes 2,8-dihydroxyadenine urolithiasis.

APRT Antibody (N-term) - References

Silva,C.H., et. al.,J. Biomol. Struct. Dyn. 25 (6), 589-597 (2008)
Di Pietro,V., et. al., Clin. Biochem. 40 (1-2), 73-80 (2007)

APRT Antibody (N-term) - Citations

- [Adenine phosphoribosyltransferase \(APRT\) deficiency: identification of a novel nonsense mutation.](#)