

**CFL1 Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2905a****Specification**

---

**CFL1 Antibody (N-term) - Product Information**

Application	<b>WB, IHC-P, FC,E</b>
Primary Accession	<a href="#">P23528</a>
Other Accession	<a href="#">P45592</a> , <a href="#">P10668</a> , <a href="#">P18760</a> , <a href="#">Q4R5C0</a> , <a href="#">Q5E9F7</a> , <a href="#">Q6B7M7</a>
Reactivity	<b>Human, Mouse</b>
Predicted	<b>Bovine, Monkey, Pig, Rat, Sheep</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>18502</b>
Antigen Region	<b>4-32</b>

**CFL1 Antibody (N-term) - Additional Information****Gene ID** 1072**Other Names**

Cofilin-1, 18 kDa phosphoprotein, p18, Cofilin, non-muscle isoform, CFL1, CFL

**Target/Specificity**

This CFL1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 4-32 amino acids from the N-terminal region of human CFL1.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CFL1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CFL1 Antibody (N-term) - Protein Information**

**Name** CFL1

**Synonyms** CFL

**Function** Binds to F-actin and exhibits pH-sensitive F-actin depolymerizing activity (PubMed:[11812157](#)). In conjunction with the subcortical maternal complex (SCMC), plays an essential role for zygotes to progress beyond the first embryonic cell divisions via regulation of actin dynamics (PubMed:[15580268](#)). Required for the centralization of the mitotic spindle and symmetric division of zygotes (By similarity). Plays a role in the regulation of cell morphology and cytoskeletal organization in epithelial cells (PubMed:[21834987](#)). Required for the up-regulation of atypical chemokine receptor ACKR2 from endosomal compartment to cell membrane, increasing its efficiency in chemokine uptake and degradation (PubMed:[23633677](#)). Required for neural tube morphogenesis and neural crest cell migration (By similarity).

**Cellular Location**

Nucleus matrix. Cytoplasm, cytoskeleton. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium {ECO:0000250|UniProtKB:P18760}. Cell projection, growth cone {ECO:0000250|UniProtKB:P18760}. Cell projection, axon {ECO:0000250|UniProtKB:P18760}. Note=Colocalizes with the actin cytoskeleton in membrane ruffles and lamellipodia. Detected at the cleavage furrow and contractile ring during cytokinesis. Almost completely in nucleus in cells exposed to heat shock or 10% dimethyl sulfoxide

**Tissue Location**


Widely distributed in various tissues.

**CFL1 Antibody (N-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**CFL1 Antibody (N-term) - Images**



Western blot analysis of CFL1 Antibody (N-term) (Cat. #AP2905a) in MCF-7, Hela cell line and mouse brain tissue lysates (35ug/lane). CFL1 (arrow) was detected using the purified Pab.(2ug/ml)



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with CFL1 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of HeLa cells using CFL1 Antibody (N-term) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### **CFL1 Antibody (N-term) - Background**

Cofilin is a widely distributed intracellular actin-modulating protein that binds and depolymerizes filamentous F-actin and inhibits the polymerization of monomeric G-actin in a pH-dependent manner.

### **CFL1 Antibody (N-term) - References**

Fazal, F., et al., J. Biol. Chem. 284 (31), 21047-21056 (2009)

### **CFL1 Antibody (N-term) - Citations**

- [Expression of UGP2 and CFL1 expression levels in benign and malignant pancreatic lesions and their clinicopathological significance.](#)