

TPI1 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2917a**Specification**

TPI1 Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P60174
Other Accession	P48500 , P00939 , Q29371 , P17751 , Q60HC9 , Q5E956
Reactivity	Human, Mouse
Predicted	Bovine, Monkey, Pig, Rabbit, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	26669
Antigen Region	68-96

TPI1 Antibody (N-term) - Additional Information**Gene ID** 7167**Other Names**

Triosephosphate isomerase, TIM, Triose-phosphate isomerase, TPI1, TPI

Target/Specificity

This TPI1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 68-96 amino acids from the N-terminal region of human TPI1.

DilutionWB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TPI1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TPI1 Antibody (N-term) - Protein Information

Name TPI1

Synonyms TPI

Function Triosephosphate isomerase is an extremely efficient metabolic enzyme that catalyzes the interconversion between dihydroxyacetone phosphate (DHAP) and D-glyceraldehyde-3-phosphate (G3P) in glycolysis and gluconeogenesis.

Cellular Location


Cytoplasm {ECO:0000255|PROSITE-ProRule:PRU10127}.

TPI1 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


TPI1 Antibody (N-term) - Images




Western blot analysis of TPI1 Antibody (N-term) (Cat. #AP2917a) in Y79(lane 1),CEM(lane 2) cell line and mouse brain tissue(lane 3) lysates (35ug/lane). TPI1 (arrow) was detected using the purified Pab.(2ug/ml)




Western blot analysis of TPI1 (arrow) using rabbit polyclonal TPI1 Antibody (N-term) (Cat. #AP2917a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the TPI1 gene.



All lanes : Anti-TPI1 Antibody (N-term) at 1:1000 dilution Lane 1: A431 whole cell lysate Lane 2: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 31 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human prostate carcinoma reacted with TPI1 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



TPI1 Antibody (N-term) (Cat. #AP2917a) flow cytometric analysis of CEM cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

TPI1 Antibody (N-term) - Background

TPI1 is an enzyme, consisting of two identical proteins, which catalyzes the isomerization of glyceraldehydes 3-phosphate (G3P) and dihydroxy-acetone phosphate (DHAP) in glycolysis and gluconeogenesis.

TPI1 Antibody (N-term) - References

Martins-de-Souza,D., et.al., BMC Psychiatry 9, 17 (2009)