

PRDX2 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2925c

Specification

PRDX2 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P32119
Other Accession	Q2PFZ3
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	21892
Antigen Region	98-127

PRDX2 Antibody (Center) - Additional Information

Gene ID 7001

Other Names

Peroxiredoxin-2, Natural killer cell-enhancing factor B, NKEF-B, PRP, Thiol-specific antioxidant protein, TSA, Thioredoxin peroxidase 1, Thioredoxin-dependent peroxide reductase 1, PRDX2, NKEFB, TDPX1

Target/Specificity

This PRDX2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 98-127 amino acids from the Central region of human PRDX2.

Dilution

WB~~1:1000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PRDX2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PRDX2 Antibody (Center) - Protein Information

Name PRDX2

Synonyms NKEFB, TDPX1

Function Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H(2)O(2).

Cellular Location


Cytoplasm.

PRDX2 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PRDX2 Antibody (Center) - Images



PRDX2 Antibody (Center) (Cat. #AP2925c) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the PRDX2 antibody detected the PRDX2 protein (arrow).



Anti-PRDX2 Antibody (Center) at 1:1000 dilution + Raji whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 22 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



Formalin-fixed and paraffin-embedded human brain tissue reacted with PRDX2 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

PRDX2 Antibody (Center) - Background

PRDX2 is a member of the peroxiredoxin family of antioxidant enzymes, which reduce hydrogen peroxide and alkyl hydroperoxides. This protein may play an antioxidant protective role in cells, and may contribute to the antiviral activity of CD8(+) T-cells. This protein may have a proliferative effect and play a role in cancer development or progression. The crystal structure of this protein has been resolved to 2.7 angstroms.

PRDX2 Antibody (Center) - References

Engstrom, K.S., et al., Mutat. Res. (2009)