

YWHAG Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2939c**Specification**

YWHAG Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P61981
Other Accession	O6NRY9 , O6PCG0 , P61983 , P61982 , Q5F3W6 , P68252 , P68253
Reactivity	Human, Mouse
Predicted	Bovine, Chicken, Rat, Sheep, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	28303
Antigen Region	136-164

YWHAG Antibody (Center) - Additional Information**Gene ID** 7532**Other Names**

14-3-3 protein gamma, Protein kinase C inhibitor protein 1, KCIP-1, 14-3-3 protein gamma, N-terminally processed, YWHAG

Target/Specificity

This YWHAG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 136-164 amino acids from the Central region of human YWHAG.

DilutionWB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

YWHAG Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

YWHAG Antibody (Center) - Protein Information

Name YWHAG ([HGNC:12852](#))

Function Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed:[15696159](#), PubMed:[16511572](#), PubMed:[36732624](#)). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed:[15696159](#), PubMed:[16511572](#), PubMed:[36732624](#)). Binding generally results in the modulation of the activity of the binding partner (PubMed:[16511572](#)). Promotes inactivation of WDR24 component of the GATOR2 complex by binding to phosphorylated WDR24 (PubMed:[36732624](#)). Participates in the positive regulation of NMDA glutamate receptor activity by promoting the L- glutamate secretion through interaction with BEST1 (PubMed:[29121962](#)).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P68252}.

Tissue Location


Highly expressed in brain, skeletal muscle, and heart.

YWHAG Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


YWHAG Antibody (Center) - Images




Western blot analysis of YWHAG Antibody (Center) (Cat. #AP2939c) in mouse cerebellum tissue lysates (35ug/lane). YWHAG (arrow) was detected using the purified Pab.




Western blot analysis of YWHAG (arrow) using rabbit polyclonal YWHAG Antibody (Center) (Cat. #AP2939c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the YWHAG gene.



Western blot analysis of lysate from HeLa cell line, using YWHAG Antibody (Center)(Cat. #AP2939c). AP2939c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



Formalin-fixed and paraffin-embedded human colon carcinoma reacted with YWHAG Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of widr cells using YWHAG Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram) FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

YWHAG Antibody (Center) - Background

YWHAG belongs to the 14-3-3 family of proteins which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 100% identical to the rat ortholog. It is induced by growth factors in human vascular smooth muscle cells, and is also highly expressed in skeletal and heart muscles, suggesting an important role for this protein in muscle tissue. It has been shown to interact with RAF1 and protein kinase C, proteins involved in various signal transduction pathways.

YWHAG Antibody (Center) - References

Jagemann, L.R., et al., J. Biol. Chem. 283 (25), 17450-17462 (2008)