

Phospho-ErbB2(Y1248) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3094a

Specification

Phospho-ErbB2(Y1248) Antibody - Product Information

Application	WB, DB,E
Primary Accession	P04626
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	137910

Phospho-ErbB2(Y1248) Antibody - Additional Information

Gene ID 2064

Other Names

Receptor tyrosine-protein kinase erbB-2, Metastatic lymph node gene 19 protein, MLN 19, Proto-oncogene Neu, Proto-oncogene c-ErbB-2, Tyrosine kinase-type cell surface receptor HER2, p185erbB2, CD340, ERBB2, HER2, MLN19, NEU, NGL

Target/Specificity

This ErbB2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y1248 of human ErbB2.

Dilution

WB~~1:8000

DB~~1:500

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is first purified by protein G affinity chromatography. Then, the antibody fraction is peptide affinity purified in a 2-step procedure with control and phosphorylated peptides. The phospho-specific antibody is eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-ErbB2(Y1248) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-ErbB2(Y1248) Antibody - Protein Information

Name ERBB2

Synonyms HER2, MLN19, NEU, NGL

Function Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell projection, ruffle membrane; Single-pass type I membrane protein. Note=Internalized from the cell membrane in response to EGF stimulation. [Isoform 2]: Cytoplasm. Nucleus.

Tissue Location


Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.

Phospho-ErbB2(Y1248) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Phospho-ErbB2(Y1248) Antibody - Images



Western blot analysis of extracts from A431 cells, untreated or treated with EGF, 100ng/ml, using phospho-ERBB2-Y1248(left) or ERBB2 antibody(right)



Western blot analysis of extracts from A431 cells, untreated or treated with EGF, 100ng/ml, using phospho-ERBB2-pY1248(L)(left) or ERBB2 antibody(right)



Dot blot analysis of anti-Phospho-ErbB2-pY1248(M) Phospho-specific Pab (Cat. #AP3094a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Phospho-ErbB2(Y1248) Antibody - Background

ErbB2 is a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors.

Phospho-ErbB2(Y1248) Antibody - References

Provinciali, M., et al., *Int. J. Cancer* 115(1):36-45 (2005). Yee, L.D., et al., *J. Nutr.* 135(5):983-988 (2005). Beckers, J., et al., *Int. J. Cancer* 114(4):590-597 (2005). Brandt, B., et al., *Biochem. Biophys.*

Res. Commun. 329(1):318-323 (2005). Ghatak, S., et al., J. Biol. Chem. 280(10):8875-8883 (2005).