

Phospho-MeCP2(S80) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3595a

Specification

Phospho-MeCP2(S80) Antibody - Product Information

Application	DB,E
Primary Accession	P51608
Other Accession	Q00566 , Q9Z2D6 , Q95LG8
Reactivity	Human
Predicted	Monkey, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52441

Phospho-MeCP2(S80) Antibody - Additional Information

Gene ID 4204

Other Names

Methyl-CpG-binding protein 2, MeCp-2 protein, MeCp2, MECP2

Target/Specificity

This MeCP2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S80 of human MeCP2.

Dilution

DB~~1:500

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-MeCP2(S80) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-MeCP2(S80) Antibody - Protein Information

Name MECP2

Function Chromosomal protein that binds to methylated DNA. It can bind specifically to a single methyl-CpG pair. It is not influenced by sequences flanking the methyl-CpGs. Mediates

transcriptional repression through interaction with histone deacetylase and the corepressor SIN3A. Binds both 5-methylcytosine (5mC) and 5-hydroxymethylcytosine (5hmC)- containing DNA, with a preference for 5-methylcytosine (5mC).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9Z2D6}. Note=Colocalized with methyl-CpG in the genome. Colocalized with TBL1X to the heterochromatin foci.

Tissue Location


Present in all adult somatic tissues tested.

Phospho-MeCP2(S80) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-MeCP2(S80) Antibody - Images



Dot blot analysis of anti-Phospho-MeCP2-pS80 Antibody (Cat.#AP3595a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Phospho-MeCP2(S80) Antibody - Background

DNA methylation is the major modification of eukaryotic genomes and plays an essential role in mammalian development. Human proteins MECP2, MBD1, MBD2, MBD3, and MBD4 comprise a family of nuclear proteins related by the presence in each of a methyl-CpG binding domain (MBD). Each of these proteins, with the exception of MBD3, is capable of binding specifically to methylated DNA. MECP2, MBD1 and MBD2 can also repress transcription from methylated gene promoters. In contrast to other MBD family members, MECP2 is X-linked and subject to X inactivation. MECP2 is dispensable in stem cells, but is essential for embryonic development. MECP2 gene mutations are

the cause of some cases of Rett syndrome, a progressive neurologic developmental disorder and one of the most common causes of mental retardation in females.

Phospho-MeCP2(S80) Antibody - References

Mnatzakanian, G.N., et al., Nat. Genet. 36(4):339-341 (2004).
Laccone, F., et al., Hum. Mutat. 23(3):234-244 (2004).
Suzuki, M., et al., Oncogene 22(54):8688-8698 (2003).
Balmer, D., et al., J. Mol. Med. 81(1):61-68 (2003).
Hagberg, B., et al., Eur. J. Paediatr. Neurol. 7(6):417-421 (2003).

Phospho-MeCP2(S80) Antibody - Citations

- [Expression of Phospho-MeCP2s in the Developing Rat Brain and Function of Postnatal MeCP2 in Cerebellar Neural Cell Development.](#)