

**Phospho-PDX1(T11) Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP3669a**

**Specification**

---

**Phospho-PDX1(T11) Antibody - Product Information**

|                   |   |
|-------------------|---|
| Application       | <b>IF, DB,E</b>   |
| Primary Accession | <a href="#">P52945</a>  |
| Other Accession   | <a href="#">P52947</a> , <a href="#">P52946</a> , <a href="#">NP_000200</a> |
| Reactivity        | <b>Human</b>  |
| Predicted         | <b>Mouse, Rat</b>   |
| Host              | <b>Rabbit</b>   |
| Clonality         | <b>Polyclonal</b>   |
| Isotype           | <b>Rabbit IgG</b>   |

**Phospho-PDX1(T11) Antibody - Additional Information**

**Gene ID** 3651

**Other Names**

Pancreas/duodenum homeobox protein 1, PDX-1, Glucose-sensitive factor, GSF, Insulin promoter factor 1, IPF-1, Insulin upstream factor 1, IUF-1, Islet/duodenum homeobox-1, IDX-1, Somatostatin-transactivating factor 1, STF-1, PDX1, IPF1, STF1

**Target/Specificity**

This PDX1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding T11 of human PDX1.

**Dilution**

IF~~1:100  
DB~~1:500

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Phospho-PDX1(T11) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Phospho-PDX1(T11) Antibody - Protein Information**

**Name** PDX1

## Synonyms IPF1, STF1

**Function** Activates insulin, somatostatin, glucokinase, islet amyloid polypeptide and glucose transporter type 2 gene transcription. Particularly involved in glucose-dependent regulation of insulin gene transcription. As part of a PDX1:PBX1b:MEIS2b complex in pancreatic acinar cells is involved in the transcriptional activation of the ELA1 enhancer; the complex binds to the enhancer B element and cooperates with the transcription factor 1 complex (PTF1) bound to the enhancer A element. Binds preferentially the DNA motif 5'-[CT]TAAT[TG]-3'. During development, specifies the early pancreatic epithelium, permitting its proliferation, branching and subsequent differentiation. At adult stage, required for maintaining the hormone-producing phenotype of the beta-cell.

## Cellular Location

Nucleus. Cytoplasm, cytosol.

## Tissue Location


Duodenum and pancreas (Langerhans islet beta cells and small subsets of endocrine non-beta-cells, at low levels in acinar cells)

## Phospho-PDX1(T11) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Phospho-PDX1(T11) Antibody - Images



Fluorescent confocal image of SY5Y cells stained with phospho-PDX1-T11 antibody. SY5Y cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with AP3669a phospho-PDX1-T11 primary antibody (1:100, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was

used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue) (10 µg/ml, 5 min). Note the highly specific localization of the phospho-PDX1 immunosignal mainly to the nucleus.



Dot blot analysis of PDX1 Antibody (T11) Pab (Cat. #AP3669a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

#### **Phospho-PDX1(T11) Antibody - Background**

PDX1 is a transcriptional activator of several genes, including insulin, somatostatin, glucokinase, islet amyloid polypeptide, and glucose transporter type 2. The encoded nuclear protein is involved in the early development of the pancreas and plays a major role in glucose-dependent regulation of insulin gene expression. Defects in this protein are a cause of pancreatic agenesis, which can lead to early-onset insulin-dependent diabetes mellitus (NIDDM), as well as maturity onset diabetes of the young type 4 (MODY4).

#### **Phospho-PDX1(T11) Antibody - References**

References for protein:

1. Nicolino, M., et al., Diabetes 59 (3), 733-740 (2010)
2. Cai, J., et al., J Mol Cell Biol 2 (1), 50-60 (2010)

References for SY5Y (SH-SY5Y; ATCC#CRL-2266): 1. Ross RA, et al. Coordinate morphological and biochemical interconversion of human neuroblastoma cells. J. Natl. Cancer Inst. 71: 741-749, 1983. [PubMed: 6137586]; 2. Biedler JL, et al. Multiple neurotransmitter synthesis by human neuroblastoma cell lines and clones. Cancer Res. 38: 3751-3757, 1978. [PubMed: 29704].