

Phospho-JMJD2B(S566) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3754a

Specification

Phospho-JMJD2B(S566) Antibody - Product Information

Application	DB,E
Primary Accession	O94953
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	121897

Phospho-JMJD2B(S566) Antibody - Additional Information

Gene ID 23030

Other Names

Lysine-specific demethylase 4B, 11411-, JmjC domain-containing histone demethylation protein 3B, Jumonji domain-containing protein 2B, KDM4B, JHDM3B, JMJD2B, KIAA0876

Target/Specificity

This JMJD2B Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S566 of human JMJD2B.

Dilution

DB~~1:500

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-JMJD2B(S566) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-JMJD2B(S566) Antibody - Protein Information

Name KDM4B

Synonyms JHDM3B, JMJD2B, KIAA0876

Function Histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby

playing a role in histone code. Does not demethylate histone H3 'Lys-4', H3 'Lys-27', H3 'Lys-36' nor H4 'Lys-20'. Only able to demethylate trimethylated H3 'Lys-9', with a weaker activity than KDM4A, KDM4C and KDM4D. Demethylation of Lys residue generates formaldehyde and succinate (PubMed:[16603238](#), PubMed:[28262558](#)). Plays a critical role in the development of the central nervous system (CNS).

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00537, ECO:0000269|PubMed:15927959}

Phospho-JMJD2B(S566) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-JMJD2B(S566) Antibody - Images



Dot blot analysis of anti-Phospho-JMJD2B-S566 Phospho-specific Pab (Cat. #AP3754a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.6ug per ml.

Phospho-JMJD2B(S566) Antibody - Background

Histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a role in histone code. Does not demethylate histone H3 'Lys-4', H3 'Lys-27', H3 'Lys-36' nor H4 'Lys-20'. Only able to demethylate trimethylated H3 'Lys-9', with a weaker activity than KDM4A, KDM4C and KDM4D. Demethylation of Lys residue generates formaldehyde and succinate.

Phospho-JMJD2B(S566) Antibody - References

Yang, J., et al. Cancer Res. 70(16):6456-6466(2010)
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :

Beyer, S., et al. J. Biol. Chem. 283(52):36542-36552(2008)
Pollard, P.J., et al. Biochem. J. 416(3):387-394(2008)
Kato, Y., et al. Int. J. Mol. Med. 20(2):269-273(2007)