

**ABCC11 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP4787a****Specification**

---

**ABCC11 Antibody (N-term) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">O96J66</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>154301</b>
Antigen Region	<b>343-372</b>

**ABCC11 Antibody (N-term) - Additional Information****Gene ID** 85320**Other Names**

ATP-binding cassette sub-family C member 11, Multidrug resistance-associated protein 8, ABCC11, MRP8

**Target/Specificity**

This ABCC11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 343-372 amino acids from the N-terminal region of human ABCC11.

**Dilution**WB~~1:1000  
IHC-P~~1:50~100**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ABCC11 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**ABCC11 Antibody (N-term) - Protein Information****Name** ABCC11 ([HGNC:14639](#))**Function** ATP-dependent transporter of the ATP-binding cassette (ABC) family that actively

extrudes physiological compounds and xenobiotics from cells. Plays a role in physiological processes involving bile acids, conjugated steroids and cyclic nucleotides, including cAMP and cGMP (PubMed:[12764137](#), PubMed:[15537867](#)). Mediates the ATP-dependent efflux of a range of physiological lipophilic anions, including the glutathione S-conjugates leukotriene C4 and dinitrophenyl S- glutathione, steroid sulfates, such as dehydroepiandrosterone 3-sulfate (DHEAS) and estrone 3-sulfate, glucuronides such as estradiol 17-beta- D-glucuronide (E(2)17betaG), the monoanionic bile acids glycocholate and taurocholate, and methotrexate (PubMed:[15537867](#), PubMed:[16359813](#), PubMed:[25896536](#)). Plays a role in the transport of earwax components (PubMed:[16444273](#), PubMed:[19383836](#)). Participates in the secretion of odorants and their precursors from the apocrine sweat glands, including the secretion of glutamine conjugates, as well as the Cys-Gly-(S) conjugates of 3-methyl-3-sulfanyl-hexanol (PubMed:[19710689](#)). Involved in the cellular extrusion of nucleotide analogs, hence conferring resistance to various drugs, including clinically relevant drugs such as 5-fluorouracil (5-FU) and methotrexate (PubMed:[12764137](#), PubMed:[15537867](#), PubMed:[25896536](#)).

#### Cellular Location

Cell membrane; Multi-pass membrane protein. Vacuole membrane Cytoplasmic vesicle membrane. Apical cell membrane; Multi-pass membrane protein

#### Tissue Location


Expressed in apocrine glands (at protein level) (PubMed:[19383836](#), PubMed:[19710689](#)). Expressed at moderate levels in breast and testis and at very low levels in liver, brain and placenta (PubMed:[11483364](#), PubMed:[11591886](#), PubMed:[16359813](#)). Localizes to axons of the central and peripheral nervous system (at protein level) (PubMed:[16359813](#)).

#### ABCC11 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### ABCC11 Antibody (N-term) - Images



Western blot analysis of ABCC11 Antibody (N-term) (Cat. #AP4787a) in WiDr cell line lysates (35ug/lane). ABCC11 (arrow) was detected using the purified Pab.



ABCC11 Antibody (N-term) (Cat. #AP4787a) IHC analysis in formalin fixed and paraffin embedded breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ABCC11 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **ABCC11 Antibody (N-term) - Background**

ABCC11 is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This ABC full transporter is a member of the MRP subfamily which is involved in multi-drug resistance. The product of this gene participates in physiological processes involving bile acids, conjugated steroids, and cyclic nucleotides. In addition, a SNP in this gene is responsible for determination of human earwax type. This gene and family member ABCC12 are determined to be derived by duplication and are both localized to chromosome 16q12.1.

#### **ABCC11 Antibody (N-term) - References**

- Martin, A., et al. J. Invest. Dermatol. 130(2):529-540(2010)
- J. Hum. Genet. 54(9):499-503(2009)
- Sato, T., et al. J. Hum. Genet. 54(7):409-413(2009)