

**GSDMA Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP4887a**

**Specification**

---

**GSDMA Antibody (N-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">O96QA5</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	49365
Antigen Region	66-94

**GSDMA Antibody (N-term) - Additional Information**

**Gene ID** 284110

**Other Names**

Gasdermin-A, Gasdermin-1, GSDMA, GSDM, GSDM1

**Target/Specificity**

This GSDMA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 66-94 amino acids from the N-terminal region of human GSDMA.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GSDMA Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**GSDMA Antibody (N-term) - Protein Information**

**Name** GSDMA ([HGNC:13311](#))

**Function** [Gasdermin-A]: This form constitutes the precursor of the pore-forming protein and acts

as a sensor of infection: upon infection by *S.pyogenes*, specifically cleaved by *S.pyogenes* effector protein SpeB in epithelial cells, releasing the N-terminal moiety (Gasdermin-A, N- terminal) that binds to membranes and forms pores, triggering pyroptosis.

#### Cellular Location

[Gasdermin-A]: Cytoplasm, perinuclear region. Cytoplasm, cytosol  
{ECO:0000250|UniProtKB:Q5Y4Y6}

#### Tissue Location


Expressed predominantly in the gastrointestinal tract and, at a lower level, in the skin. Also detected in mammary gland. In the gastrointestinal tract, mainly expressed in differentiated cells, including the differentiated cell layer of esophagus and mucus-secreting pit cells of the gastric epithelium Down-regulated in gastric cancer cells.

### GSDMA Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### GSDMA Antibody (N-term) - Images



Western blot analysis of GSDMA Antibody (N-term) (Cat. #AP4887a) in mouse liver tissue lysates (35ug/lane). GSDMA (arrow) was detected using the purified Pab.



GSDMA Antibody (N-term) (Cat. #AP4887a) IHC analysis in formalin fixed and paraffin embedded human skin carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the GSDMA Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



GSDMA Antibody (N-term) (Cat. #AP4887a) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

**GSDMA Antibody (N-term) - Background**

GSDMA induces apoptosis.

**GSDMA Antibody (N-term) - References**

Soranzo, N., et al. Nat. Genet. 41(11):1182-1190(2009)  
 Saeki, N., et al. Genes Chromosomes Cancer 48(3):261-271(2009)  
 Tamura, M., et al. Genomics 89(5):618-629(2007)