

## Cyclin D2 Antibody (Ab-280)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP50014

## **Specification**

## Cyclin D2 Antibody (Ab-280) - Product Information

Application WB
Primary Accession P30279

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 33 KDa
Antigen Region 253-285

## Cyclin D2 Antibody (Ab-280) - Additional Information

Gene ID 894

**Other Names** 

G1/S-specific cyclin-D2, CCND2

**Dilution** 

WB~~ 1:500-1:1000

#### **Format**

Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

**Storage Conditions** -20°C

## Cyclin D2 Antibody (Ab-280) - Protein Information

Name CCND2 {ECO:0000303|PubMed:1386336, ECO:0000312|HGNC:HGNC:1583}

### **Function**

Regulatory component of the cyclin D2-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition (PubMed:<a href="http://www.uniprot.org/citations/18827403" target="\_blank">18827403" target="\_blank">18827403" target="\_blank">18827403</a>, PubMed:<a href="http://www.uniprot.org/citations/8114739" target="\_blank">8114739</a>). Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase (PubMed:<a

href="http://www.uniprot.org/citations/18827403" target="\_blank">18827403</a>, PubMed:<a href="http://www.uniprot.org/citations/8114739" target=" blank">8114739</a>).

Hypophosphorylates RB1 in early G(1) phase (PubMed:<a

href="http://www.uniprot.org/citations/18827403" target="\_blank">18827403</a>, PubMed:<a href="http://www.uniprot.org/citations/8114739" target=" blank">8114739</a>). Cyclin D-CDK4





complexes are major integrators of various mitogenenic and antimitogenic signals (PubMed:<a href="http://www.uniprot.org/citations/18827403" target="\_blank">18827403</a>, PubMed:<a href="http://www.uniprot.org/citations/8114739" target=" blank">8114739</a>).

#### **Cellular Location**


Nucleus. Cytoplasm. Nucleus membrane. Note=Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated into the nucleus through interaction with KIP/CIP family members

## Cyclin D2 Antibody (Ab-280) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## Cyclin D2 Antibody (Ab-280) - Images



Western blot analysis of lysate from RD cell line, using Cyclin D2 Antibody (Ab-280)(B8336). B8336 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

# Cyclin D2 Antibody (Ab-280) - Background

Regulatory component of the cyclin D2-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenenic and antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and





Tel: 858.875.1900 Fax: 858.875.1999

repressing its transcriptional activity. Component of the ternary complex, cyclin D2/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (By similarity).

## Cyclin D2 Antibody (Ab-280) - References

Xiong Y., et al. Genomics 13:575-584(1992). Palmero I., et al. Oncogene 8:1049-1054(1993). Miyajima N., et al. Submitted (MAR-1993) to the EMBL/GenBank/DDBJ databases. Kalnine N., et al. Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases. Ota T., et al. Nat. Genet. 36:40-45(2004).