

**hnRNP Q Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP50037****Specification**

---

**hnRNP Q Antibody - Product Information**

Application	IF, WB
Primary Accession	<a href="#">O60506</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70,66,63,59,46 KDa
Antigen Region	255-284

**hnRNP Q Antibody - Additional Information****Gene ID** 10492**Other Names**

Heterogeneous nuclear ribonucleoprotein Q, hnRNP Q, Glycine- and tyrosine-rich RNA-binding protein, GRY-RBP, NS1-associated protein 1, Synaptotagmin-binding, cytoplasmic RNA-interacting protein, SYNCRIP, HNRPQ, NSAP1

**Dilution**

IF~~1:100

WB~~ 1:1000

**Format**

Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

**Storage Conditions**

-20°C

**hnRNP Q Antibody - Protein Information****Name** SYNCRIP**Synonyms** HNRPQ, NSAP1**Function**

Heterogenous nuclear ribonucleoprotein (hnRNP) implicated in mRNA processing mechanisms. Component of the CRD-mediated complex that promotes MYC mRNA stability. Isoform 1, isoform 2 and isoform 3 are associated in vitro with pre-mRNA, splicing intermediates and mature mRNA protein complexes. Isoform 1 binds to apoB mRNA AU-rich sequences. Isoform 1 is part of the APOB mRNA editosome complex and may modulate the postranscriptional C to U RNA-editing of the APOB mRNA through either by binding to A1CF (APOBEC1 complementation factor), to APOBEC1 or to RNA itself. May be involved in translationally coupled mRNA turnover. Implicated

with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. Interacts in vitro preferentially with poly(A) and poly(U) RNA sequences. Isoform 3 may be involved in cytoplasmic vesicle-based mRNA transport through interaction with synaptotagmins. Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes. Upon interferon-gamma activation assembles into the GAIT complex which binds to stem loop- containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation; seems not to be essential for GAIT complex function.

#### Cellular Location

Cytoplasm. Microsome {ECO:0000250|UniProtKB:Q7TMK9} Endoplasmic reticulum. Nucleus {ECO:0000250|UniProtKB:Q7TMK9}. Note=The tyrosine phosphorylated form bound to RNA is found in microsomes (By similarity). Localized in cytoplasmic mRNP granules containing untranslated mRNAs (By similarity). {ECO:0000250|UniProtKB:O43390, ECO:0000250|UniProtKB:Q7TMK9} [Isoform 2]: Nucleus, nucleoplasm {ECO:0000250|UniProtKB:Q7TMK9}. Note=Expressed predominantly in the nucleoplasm. {ECO:0000250|UniProtKB:Q7TMK9}

#### Tissue Location


Ubiquitously expressed. Detected in heart, brain, pancreas, placenta, spleen, lung, liver, skeletal muscle, kidney, thymus, prostate, uterus, small intestine, colon, peripheral blood and testis.

#### hnRNP Q Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### hnRNP Q Antibody - Images



Immunofluorescence analysis of HepG2 cells, using hnRNP Q antibody.



Western blot analysis of lysates from HT-29, HeLa cell line (from left to right), using hnRNP Q Antibody (C10139). C10139 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 µg per lane.

### hnRNP Q Antibody - Background

Heterogeneous nuclear ribonucleoprotein (hnRNP) implicated in mRNA processing mechanisms. Component of the CRD-mediated complex that promotes MYC mRNA stability. Isoform 1, isoform 2 and isoform 3 are associated in vitro with pre-mRNA, splicing intermediates and mature mRNA protein complexes. Isoform 1 binds to apoB mRNA AU-rich sequences. Isoform 1 is part of the APOB mRNA editosome complex and may modulate the posttranscriptional C to U RNA-editing of the APOB mRNA through either by binding to A1CF (APOBEC1 complementation factor), to APOBEC1 or to RNA itself. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. Interacts in vitro preferentially with poly(A) and poly(U) RNA sequences. Isoform 3 may be involved in cytoplasmic vesicle-based mRNA transport through interaction with synaptotagmins. Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes. Upon interferon-gamma activation assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruloplasmin) and suppresses their translation; seems not to be essential for GAIT complex function.

### hnRNP Q Antibody - References

- Harris C.E., et al. *J. Virol.* 73:72-80(1999).
- Du G., et al. *Chin. Sci. Bull.* 45:343-349(2000).
- Mourelatos Z., et al. *EMBO J.* 20:5443-5452(2001).
- Suzuki Y., et al. Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
- Mungall A.J., et al. *Nature* 425:805-811(2003).