

MAD2L1 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP50070**Specification**

MAD2L1 Antibody - Product Information

Application	WB
Primary Accession	O13257
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24,10 KDa
Antigen Region	95-126

MAD2L1 Antibody - Additional Information**Gene ID** 4085**Other Names**

Mitotic spindle assembly checkpoint protein MAD2A, HsMAD2, Mitotic arrest deficient 2-like protein 1, MAD2-like protein 1, MAD2L1, MAD2

Dilution

WB~~ 1:1000

Format

Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions

-20°C

MAD2L1 Antibody - Protein Information**Name** MAD2L1**Synonyms** MAD2**Function**

Component of the spindle-assembly checkpoint that prevents the onset of anaphase until all chromosomes are properly aligned at the metaphase plate (PubMed: [15024386](http://www.uniprot.org/citations/15024386), PubMed: [29162720](http://www.uniprot.org/citations/29162720)). In the closed conformation (C-MAD2) forms a heterotetrameric complex with MAD1L1 at unattached kinetochores during prometaphase, the complex recruits open conformation molecules of MAD2L1 (O-MAD2) and then promotes the conversion of O-MAD2 to C-MAD2 (PubMed: [29162720](http://www.uniprot.org/citations/29162720)). Required for the execution of the mitotic checkpoint which monitors the process of kinetochore-spindle

attachment and inhibits the activity of the anaphase promoting complex by sequestering CDC20 until all chromosomes are aligned at the metaphase plate (PubMed:10700282, PubMed:11804586, PubMed:15024386).

Cellular Location


Nucleus. Chromosome, centromere, kinetochore. Cytoplasm. Cytoplasm, cytoskeleton, spindle pole Note=Recruited by MAD1L1 to unattached kinetochores (Probable) Recruited to the nuclear pore complex by TPR during interphase Recruited to kinetochores in late prometaphase after BUB1, CENPF, BUB1B and CENPE. Kinetochore association requires the presence of NEK2 Kinetochore association is repressed by UBD. Sequestered to the cytoplasm upon interaction with isoform 3 of MAD1L1 (PubMed:19010891) {ECO:0000269|PubMed:19010891, ECO:0000305}

MAD2L1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MAD2L1 Antibody - Images



Western blot analysis of lysate from HT-29 cell line, using MAD2L1 Antibody (C21657). C21657 was diluted at 1:1000. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

MAD2L1 Antibody - Background

Component of the spindle-assembly checkpoint that prevents the onset of anaphase until all chromosomes are properly aligned at the metaphase plate. Required for the execution of the

mitotic checkpoint which monitors the process of kinetochore- spindle attachment and inhibits the activity of the anaphase promoting complex by sequestering CDC20 until all chromosomes are aligned at the metaphase plate.

MAD2L1 Antibody - References

Li Y.,et al.Science 274:246-248(1996).

Gemma A.,et al.Lung Cancer 32:289-295(2001).

Jin D.-Y.,et al.Submitted (JUL-1995) to the EMBL/GenBank/DDBJ databases.

Klebert S.,et al.Submitted (OCT-1997) to the EMBL/GenBank/DDBJ databases.

Nobori T.,et al.Submitted (FEB-2001) to the EMBL/GenBank/DDBJ databases.